

Advanced Lectures in Mathematics (ALM)

- ALM 1: Superstring Theory
- ALM 2: Asymptotic Theory in Probability and Statistics with Applications
- ALM 3: Computational Conformal Geometry
- ALM 4: Variational Principles for Discrete Surfaces
- ALM 6: Geometry, Analysis and Topology of Discrete Groups
- ALM 7: Handbook of Geometric Analysis, No. 1
- ALM 8: Recent Developments in Algebra and Related Areas
- ALM 9: Automorphic Forms and the Langlands Program
- ALM 10: Trends in Partial Differential Equations
- ALM 11: Recent Advances in Geometric Analysis
- ALM 12: Cohomology of Groups and Algebraic K-theory
- ALM 13: Handbook of Geometric Analysis, No. 2
- ALM 14: Handbook of Geometric Analysis, No. 3
- ALM 15: An Introduction to Groups and Lattices: Finite Groups and Positive Definite Rational Lattices
- ALM 16: Transformation Groups and Moduli Spaces of Curves
- ALM 17: Geometry and Analysis, No. 1
- ALM 18: Geometry and Analysis, No. 2
- ALM 19: Arithmetic Geometry and Automorphic Forms
- ALM 20: Surveys in Geometric Analysis and Relativity
- ALM 21: Advances in Geometric Analysis
- ALM 22: Differential Geometry: Under the Influence of S.-S. Chern
- ALM 23: Recent Developments in Geometry and Analysis
- ALM 24: Handbook of Moduli, Volume I
- ALM 25: Handbook of Moduli, Volume II
- ALM 26: Handbook of Moduli, Volume III

Advanced Lectures in Mathematics
Volume XXIV

Handbook of Moduli

Volume I

edited by

Gavril Farkas · Ian Morrison

 International Press
www.intlpress.com

 高等教育出版社
HIGHER EDUCATION PRESS

Advanced Lectures in Mathematics, Volume XXIV
Handbook of Moduli, Volume I

Volume Editors:

Gavril Farkas (Humboldt-Universität, Berlin)

Ian Morrison (Fordham University, New York)

2010 Mathematics Subject Classification. Primary: 14D20. Secondary: 14H10,
14K10, 11G15, 14N35, 14E30, 14D15.

Copyright © 2013 by International Press, Somerville, Massachusetts, U.S.A., and by
Higher Education Press, Beijing, China.

This work is published and sold in China exclusively by Higher Education Press
of China.

All rights reserved. Individual readers of this publication, and non-profit libraries acting for them, are permitted to make fair use of the material, such as to copy a chapter for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgement of the source is given. Republication, systematic copying, or mass reproduction of any material in this publication is permitted only under license from International Press. Excluded from these provisions is material in articles to which the author holds the copyright. (If the author holds copyright, notice of this will be given with the article.) In such cases, requests for permission to use or reprint should be addressed directly to the author.

ISBN: 978-1-57146-257-2

Printed in the United States of America.

17 16 15 14 13 1 2 3 4 5 6 7 8 9

ADVANCED LECTURES IN MATHEMATICS

Executive Editors

Shing-Tung Yau
Harvard University
Cambridge, Mass., U.S.A.

Kefeng Liu
University of California at Los Angeles
Los Angeles, Calif., U.S.A.

Lizhen Ji
University of Michigan, Ann Arbor
Ann Arbor, Mich., U.S.A.

Editorial Board

Chongqing Cheng
Nanjing University
Nanjing, China

Tatsien Li
Fudan University
Shanghai, China

Zhong-Ci Shi
Institute of Computational Mathematics
Chinese Academy of Sciences (CAS)
Beijing, China

Zhiying Wen
Tsinghua University
Beijing, China

Zhouping Xin
The Chinese University of Hong Kong
Hong Kong, China

Lo Yang
Institute of Mathematics
Chinese Academy of Sciences (CAS)
Beijing, China

Weiping Zhang
Nankai University
Tianjin, China

Xiping Zhu
Sun Yat-sen University
Guangzhou, China

Xiangyu Zhou
Institute of Mathematics
Chinese Academy of Sciences (CAS)
Beijing, China

The Handbook of Moduli is dedicated to the memory of Eckart Viehweg, whose untimely death precluded a planned contribution, and to David Mumford, who first proposed the project, for all that they both did to nurture its subject; and to Angela Ortega and Jane Reynolds for everything that they do to sustain its editors.

Contents

Volume I

Preface	
<i>Gavril Farkas and Ian Morrison</i>	xiii
Logarithmic geometry and moduli	
<i>Dan Abramovich, Qile Chen, Danny Gillam, Yuhao Huang, Martin Olsson, Matthew Satriano and Shenghao Sun</i>	1
Invariant Hilbert schemes	
<i>Michel Brion</i>	63
Algebraic and tropical curves: comparing their moduli spaces	
<i>Lucia Caporaso</i>	119
A superficial working guide to deformations and moduli	
<i>F. Catanese</i>	161
Moduli spaces of hyperbolic surfaces and their Weil–Petersson volumes	
<i>Norman Do</i>	217
Equivariant geometry and the cohomology of the moduli space of curves	
<i>Dan Edidin</i>	259
Tautological and non-tautological cohomology of the moduli space of curves	
<i>C. Faber and R. Pandharipande</i>	293
Alternate compactifications of moduli spaces of curves	
<i>Maksym Fedorchuk and David Ishii Smyth</i>	331
The cohomology of the moduli space of Abelian varieties	
<i>Gerard van der Geer</i>	415
Moduli of K3 surfaces and irreducible symplectic manifolds	
<i>V. Gritsenko, K. Hulek and G.K. Sankaran</i>	459
Normal functions and the geometry of moduli spaces of curves	
<i>Richard Hain</i>	527

Volume II

Parameter spaces of curves <i>Joe Harris</i>	1
Global topology of the Hitchin system <i>Tamás Hausel</i>	29
Differential forms on singular spaces, the minimal model program, and hyperbolicity of moduli stacks <i>Stefan Kebekus</i>	71
Contractible extremal rays on $\overline{M}_{0,n}$ <i>Seán Keel and James M^cKernan</i>	115
Moduli of varieties of general type <i>János Kollár</i>	131
Singularities of stable varieties <i>Sándor J Kovács</i>	159
Soliton equations and the Riemann-Schottky problem <i>I. Krichever and T. Shiota</i>	205
GIT and moduli with a twist <i>Radu Laza</i>	259
Good degenerations of moduli spaces <i>Jun Li</i>	299
Localization in Gromov-Witten theory and Orbifold Gromov-Witten theory <i>Chiu-Chu Melissa Liu</i>	353
From WZW models to modular functors <i>Eduard Looijenga</i>	427
Shimura varieties and moduli <i>J.S. Milne</i>	467
The Torelli locus and special subvarieties <i>Ben Moonen and Frans Oort</i>	549

Volume III

Birational geometry for nilpotent orbits <i>Yoshinori Namikawa</i>	1
Cell decompositions of moduli space, lattice points and Hurwitz problems <i>Paul Norbury</i>	39
Moduli of abelian varieties in mixed and in positive characteristic <i>Frans Oort</i>	75
Local models of Shimura varieties, I. Geometry and combinatorics <i>Georgios Pappas, Michael Rapoport and Brian Smithling</i>	135
Generalized theta linear series on moduli spaces of vector bundles on curves <i>Mihnea Popa</i>	219
Computer aided unirationality proofs of moduli spaces <i>Frank-Olaf Schreyer</i>	257
Deformation theory from the point of view of fibered categories <i>Mattia Talpo and Angelo Vistoli</i>	281
Mumford's conjecture — a topological outlook <i>Ulrike Tillmann</i>	399
Rational parametrizations of moduli spaces of curves <i>Alessandro Verra</i>	431
Hodge loci <i>Claire Voisin</i>	507
Homological stability for mapping class groups of surfaces <i>Nathalie Wahl</i>	547

Preface

Gavril Farkas and Ian Morrison

The title of these volumes might lead unwary readers to expect an encyclopedic survey for experts in the study of moduli problems in algebraic geometry. What they will discover is rather different. Our aims here are, first, to clarify the audience that we hope the Handbook *will* serve and the approach it *does* take to its subject and, second, to thank all those who have assisted us in helping it realize these aims.

To begin with, a bit of history. The idea for a Handbook of Moduli originated in a discussion between David Mumford and Lizhen Ji at Michigan in 2006. Lizhen and David produced a draft table of contents that was circulated at the Symposium marking David's retirement from Brown in 2007. The Handbook was originally to have been edited by Ching-Li Chai and Amnon Neeman, but the demands of their work with Takahiro Shiota as editors of the second volume of Mumford's Collected Papers took priority and, at their urging, we agreed to take over editorship in the spring of 2009.

We quickly reached the conclusion that what was needed for many topics was not a discussion of the latest results aimed at specialists, but a survey aimed at a broad community of producers (and even some consumers from cognate areas) of algebraic geometry, most of whom had little prior familiarity with the area. Our goal became a Handbook that would introduce the techniques, examples and results essential to each topic, and say enough about recent developments to prepare the reader to tackle the primary literature in the area. We particularly sought to elicit contributions that illustrated "secret handshakes", yogas and heuristics that experts use privately to guide intuition or simplify calculation but that are replaced by more formal arguments, or simply do not appear, in articles aimed at other specialists.

For many topics, the Handbook succeeds much better than we dared to hope. The credit is due entirely to the hard work of the Handbook's many authors in producing articles that conformed to the goals we had set. Again and again, we were delighted to find that authors, instead of taking the easy course of cutting and pasting from earlier surveys and primary references, had made the substantially greater effort to write the original treatments needed to bridge gaps in the literature and make important problems accessible to a wide audience for the first time.

We expect that they will reap a just reward and that their articles will be widely read and referenced. Here we want to offer them not only our sincerest thanks, but also those of the Handbook's readers, for their exceptional generosity. Many

Handbook articles were also improved by extensive and thoughtful referees' reports. We are grateful for all work that the referees did to improve the Handbook and take this opportunity to thank them collectively on behalf of the contributors.

We must, however, disclaim that the Handbook's coverage is often incomplete, in extreme cases, non-existent. The blame for these gaps is mostly ours. When we solicited contributions to the Handbook, each invitation was accompanied by a suggested topic, and we selected contributors who we thought would be able to cover their topics in the spirit discussed above. The results reflect both our knowledge and taste—of topics and of experts in them—and also, in some cases, our ignorance.

In some areas, we found it easy to produce candidate contributor–topic pairs, and to recruit the contributors we had identified. The Handbook's discussion of, for example, moduli spaces of curves is, therefore, particularly complete—some will say, not without a certain justice, excessive.

In other areas, we had more difficulty both in identifying and in enlisting candidates. A few of the more obvious gaps arose when authors who had accepted our invitation backed out after it was too late to find replacements. A more deeply felt loss—one that impacts the whole subject of moduli—was the untimely death of Eckart Viehweg, who had been one of the first to agree to contribute.

We also omitted a few topics as a courtesy to the authors of monographs devoted to them that we knew to be in preparation, others because papers treating them in the spirit we were seeking had recently appeared, and yet others because we felt that they were developing so rapidly that any contribution dealing with them would have a limited shelf-life. In hindsight, not all of these decisions were well taken.

As a result, the Handbook's treatment of moduli has some major lacunae (mirror symmetry, wall crossing formulae) and there are other topics (moduli of sheaves and bundles) which are discussed but not in the depth that their importance merits. We apologize to readers who may have hoped to find more about these subjects in the Handbook, and (with Lizhen's encouragement) we challenge experts who feel that their areas deserve a fuller exposition to offer him proposals for additional Handbook volumes devoted to them.

The Handbook also benefitted from the efforts of many other colleagues. Amnon Neeman showed considerable doggedness in recruiting us to succeed him and Ching-Li as editors. Scott Wolpert provided valuable advice on the cat-herding elements of the editor's job. Dave Bayer helped enormously in setting up the final production process both to automate complex and error prone operations and to prevent inconsistencies between the \LaTeX installations on our home systems and those at Higher Education Press.

Brian Bianchini, International Press' book production manager, ensured that we had the resources we needed throughout the Handbook's growth from the single volume originally projected to the present three. The Advanced Mathematics series editor, Lizhen Ji, was always ready to answer our questions, help with practical

difficulties, and adjust his schedule for the series to adapt to changes in ours. Liping Wang and her production staff at the Higher Education Press were unfailingly accommodating and helpful to us in resolving \LaTeX issues—even re \LaTeX ing several submissions to bring them into conformity with the Handbook style—and made every effort to ensure that the appearance of the Handbook volumes was up to the standard of their contents.

To all of them, and to many others who provided more informal help, we here offer our sincerest thanks.

Humboldt Universität, Institut für Mathematik, Unter den Linden 6, 10099 Berlin

E-mail address: farkas@mathematik.hu-berlin.de

Department of Mathematics, Fordham University, Bronx, NY 10458

E-mail address: morrison@fordham.edu

Moduli of K3 surfaces and irreducible symplectic manifolds

V. Gritsenko, K. Hulek, and G.K. Sankaran

Abstract. The name “K3 surfaces” was coined by A. Weil in 1957 when he formulated a research programme for these surfaces and their moduli. Since then, irreducible holomorphic symplectic manifolds have been introduced as a higher dimensional analogue of K3 surfaces. In this paper we present a review of this theory starting from the definition of K3 surfaces and going as far as the global Torelli theorem for irreducible holomorphic symplectic manifolds as recently proved by M. Verbitsky.

For many years the last open question of Weil’s programme was that of the geometric type of the moduli spaces of polarised K3 surfaces. We explain how this problem has been solved. Our method uses algebraic geometry, modular forms and Borcherds automorphic products. We collect and discuss the relevant facts from the theory of modular forms with respect to the orthogonal group $O(2, n)$. We also give a detailed description of quasi pull-back of automorphic Borcherds products. This part contains previously unpublished results.

Contents

1	Introduction	460
2	Periods of K3 surfaces and the Torelli theorem	463
	2.1 Lattices	463
	2.2 Markings and the period map	464
	2.3 The Torelli theorem	464
	2.4 The universal family of marked K3 surfaces	466
	2.5 Moduli spaces of polarised K3 surfaces	468
3	Irreducible symplectic manifolds	472
	3.1 Basic theory of irreducible symplectic manifolds	472
	3.2 Hodge theoretic Torelli theorem	475
	3.3 Moduli spaces of polarised irreducible symplectic manifolds	477
4	Projective models	480
5	Compactifications	482
	5.1 Modular varieties of orthogonal type	483
	5.2 The Baily-Borel compactification	485
	5.3 Toroidal compactifications	487
	5.4 Canonical singularities	490

2000 *Mathematics Subject Classification.* Primary 14J10; Secondary 14D99.

Key words and phrases. moduli, K3 surfaces, hyperkähler, modular form.

5.5	Singularities of modular varieties	492
5.6	Singularities in the interior	492
5.7	Singularities at the cusps	494
6	Modular forms and Kodaira dimension	495
6.1	Modular forms of orthogonal type	495
6.2	Rational quadratic divisors	497
6.3	Low weight cusp form trick	497
6.4	Reflective modular forms	499
7	Orthogonal groups and reflections	501
8	The quasi pull-back of modular forms	505
8.1	Quasi pull-back	505
8.2	Tube domain realisation	508
8.3	Fourier expansion at 0-dimensional cusps	508
8.4	Properties of quasi pull-back	510
9	Arithmetic of root systems	514
9.1	Vectors in E_8 and E_7	514
9.2	Binary quadratic forms in E_8	516

1. Introduction

The history of K3 surfaces goes back to classical algebraic geometry. In modern complex algebraic geometry a *K3 surface* is a compact complex surface S whose canonical bundle is trivial, i.e. $K_S \cong \mathcal{O}_S$, and whose irregularity $q(S) = h^1(S, \mathcal{O}_S) = 0$. These two facts immediately imply that $H^0(S, \Omega_S^1) = H^0(S, T_S) = H^2(S, T_S) = 0$.

The easiest examples of algebraic K3 surfaces are smooth quartic surfaces in \mathbb{P}^3 . Further examples are complete intersections of type $(2,3)$ in \mathbb{P}^4 and of type $(2,2,2)$ in \mathbb{P}^5 . Another classical example is the *Kummer surfaces*, i.e. the (desingularised) quotient of a 2-dimensional torus A by the involution $\iota: x \mapsto -x$.

The modern name “K3 surface” was coined by A. Weil in his famous “Final report on research contract AF 18(603)-57” [131, Vol II, pp 390–395]. In his comments on this report [131, Vol II, p 546] Weil writes: “Dans la seconde partie de mon rapport, il s’agit des variétés kählériennes dites K3, ainsi nommées en l’honneur de Kummer, Kodaira, Kähler et de la belle montagne K2 au Cachemire.”

In that report the following conjectures, due to Andreotti and Weil, were stated:

- (i) K3 surfaces form one family;
- (ii) all K3 surfaces are Kähler;
- (iii) the period map is surjective;
- (iv) a form of global Torelli theorem holds.

Weil also remarked that the structure of the moduli space of (polarised) K3 surfaces must be closely related to the theory of automorphic forms.

By now all of these questions have been answered positively and much progress has been made in understanding the moduli spaces of K3 surfaces. Conjecture (i) was proved by Kodaira [71, Part I, theorem 19]. Conjecture (ii) was first shown by Siu [122, Section 14]. Nowadays it can be derived from the more general theorem, due to Buchdahl and Lamari, that a compact complex surface is Kähler if and only if its first Betti number is even ([6, Theorem IV.3.1], [19],[78]). Surjectivity of the period map (conjecture (iii)) was proved for special K3 surfaces in various papers by Shah [117], [118], [119] and by Horikawa [61]. Kulikov [77] gave a proof for projective K3 surfaces, with some points clarified by Persson and Pinkham [105]. The general result for Kähler K3 surfaces was proved by Todorov [126] and Looijenga [80]. The strong Torelli theorem (i.e. conjecture (iv)) for algebraic K3 surfaces was first proved by Piatetskii-Shapiro and Shafarevich [108] with amendments by Rapoport and Shioda. It was proved for Kähler K3 surfaces (and hence for all) by Burns and Rapoport [20]. A detailed and supplemented account of the original proof was written by Looijenga and Peters [81]. Finally, Friedman [34] gave a proof using degenerations.

It should be noted, though, that Weil's definition of K3 surface was different from the standard definition used nowadays. For him a surface was K3 if its underlying differentiable structure was that of a quartic surface in \mathbb{P}^3 . Using results from Seiberg-Witten theory one can indeed show that any compact complex surface diffeomorphic to a quartic is a K3 surface.

The crucial ingredient is the fact that the plurigenera of a compact complex surface are invariant under orientation preserving diffeomorphism: see [6, Theorem IX.9.4 and Theorem IX.9.5]. This was formulated as a question in [103, p. 244] and became known as the Van de Ven conjecture. A first (partial) proof of this was published by Brussee [18] (see also [35]), and an elegant argument using Seiberg-Witten invariants can be found in Dür's thesis [29]. It follows that the only surfaces which admit an orientation preserving diffeomorphism to a quartic are K3 surfaces or possibly tori, but the latter are excluded because they are not simply connected. If a surface is diffeomorphic to a quartic surface, but with an orientation changing diffeomorphism, then its signature is -16 and its Euler number equals $c_2(S) = 24$. By the Thom-Hirzebruch index theorem $c_1^2(S) = 96$, but that contradicts the Miyaoka-Yau inequality, so such a surface cannot exist.

Moduli spaces of polarised K3 surfaces are a historically old subject, studied by the classical Italian geometers (starting with the spaces of double sextics and plane quartics in \mathbb{P}^3). Mukai extended the classical constructions and unirationality results for the moduli spaces \mathcal{F}_{2d} parametrising polarised K3 surfaces of degree $2d$ to many more cases, going as high as $d = 19$. See [89], [90], [91], [92] and [93] for details. Kondo [74] proved that the moduli spaces \mathcal{F}_{2p^2} are of general type for sufficiently large prime numbers p , but without giving an effective bound. Finally, it was shown in [46] that the moduli spaces \mathcal{F}_{2d} are of general type for $d > 61$ and

$d = 46, 50, 54, 57, 58, 60$, and later it was noticed by A. Peterson [106] that this proof also works for $d = 52$: see Theorem 6.1 below. For an account of these results see also Voisin's Bourbaki exposé [129].

In higher dimension K3 surfaces can be generalised in two ways, namely to Calabi-Yau varieties or to irreducible symplectic manifolds (or hyperkähler manifolds). In fact, together with tori, these three classes of manifolds are the building blocks of all compact Kähler manifolds with trivial first Chern class (over \mathbb{R}), also called Ricci flat manifolds: every such manifold admits a finite étale cover which is a product of tori, Calabi-Yau manifolds and irreducible symplectic manifolds (see [8], [9]).

The first examples of irreducible symplectic manifolds, by now regarded as classical, were studied by Beauville in [8], namely the Hilbert schemes of points on K3 surfaces and generalised Kummer varieties (and their deformations). Two further classes of examples were later found by O'Grady [99], [100]. The theory of irreducible symplectic manifolds, which started from work of Bogomolov, Beauville and Fujiki, was significantly advanced by Huybrechts [63] who, among other results, proved surjectivity of the period map. It was noticed early on by Debarre [24] and Namikawa [95] that the obvious generalisations of the Torelli theorem are false. Nevertheless, one can use the period map to exhibit moduli spaces of polarised irreducible symplectic manifolds as dominant finite to one covers of quotients of type IV domains by arithmetic groups. This was used in [50] and [51] to obtain general type results for many of these moduli spaces (Theorem 6.2 and Theorem 6.3 below).

Very recently Verbitsky [127] has announced a Torelli theorem for irreducible symplectic manifolds. The consequences of Verbitsky's result for the moduli problem of polarised irreducible symplectic manifolds were worked out in detail by Markman [86]. We also refer the reader to Huybrecht's Bourbaki talk [64].

The theory of K3 surfaces and irreducible symplectic manifolds is a fascinating and vast subject. We have started this introduction by giving the definition of K3 surfaces in complex geometry (which also allows for non-algebraic surfaces). The notion of K3 surface also makes perfect sense in algebraic geometry over arbitrary fields: a K3 surface is an irreducible smooth algebraic surface S defined over a field k with trivial canonical bundle and irregularity $q = h^1(S, \mathcal{O}_S) = 0$. In this article we shall, however, concentrate on the theory over the complex numbers, and especially on moduli problems. We are fully aware that we thus exclude a large area of interesting questions. Concerning K3 surfaces in positive characteristic we refer the reader to the papers by Artin [2], Ogus [102] and Nygaard [98] for a first introduction. Another aspect, which we will not touch upon, is the arithmetic of K3 surfaces. An introduction to this field can be found in Schütt's survey paper [116].

In this article we mainly survey known results. Theorem 8.18 is new, however (special cases occur in the literature) and so is Proposition 9.5.

2. Periods of K3 surfaces and the Torelli theorem

In this section we will discuss the period domain and the period map for K3 surfaces, and the local and global Torelli theorems. The case of K3 surfaces presents several special features and the existence of a straightforward global Torelli theorem is one of them.

2.1. Lattices

Let S be a K3 surface. By Noether’s formula $b_2(S) = 22$, and since S is simply connected (being diffeomorphic to a quartic hypersurface), $H^2(S, \mathbb{Z})$ is a free \mathbb{Z} -module of rank 22. The intersection form defines a non-degenerate symmetric bilinear form on $H^2(S, \mathbb{Z})$, giving it the structure of a lattice, which has signature $(3, 19)$ by the Hodge index theorem. As the intersection form is even and unimodular, this lattice is, independently of the surface S , the *K3 lattice*

$$(2.1) \quad L_{K3} := 3U \oplus 2E_8(-1)$$

where U is the hyperbolic plane (the unique even unimodular lattice of signature $(1, 1)$) and E_8 is the unique even unimodular positive definite rank 8 lattice. For any lattice L and $m \in \mathbb{Z}$, the notation $L(m)$ indicates that the quadratic form is multiplied by m , so $E_8(-1)$ is negative definite.

It was proven by Siu [122] that every K3 surface is Kähler. Today, it is known that a compact complex surface is Kähler if and only if its first Betti number is even [6, Theorem IV.3.1], which immediately implies Siu’s result. Hence $H^2(S, \mathbb{C})$ has a Hodge decomposition

$$H^2(S, \mathbb{C}) = H^{2,0}(S) \oplus H^{1,1}(S) \oplus H^{0,2}(S).$$

Since $K_S \cong \mathcal{O}_S$ it follows that $H^{2,0}(S) = H^0(S, K_S)$ is 1-dimensional and thus $H^{1,1}(S)$ has dimension 20.

Since the second cohomology has no torsion we can, via the universal coefficient theorem, consider $H^2(S, \mathbb{Z})$ as a lattice in $H^2(S, \mathbb{C})$. The *Néron-Severi group* is the intersection $NS(S) = H^2(S, \mathbb{Z}) \cap H^{1,1}(S)$, which, in this case, coincides with the Picard group $\text{Pic}(S)$. The *transcendental lattice* is defined as the smallest lattice whose complexification contains a generator ω of $H^0(S, K_S)$. If the intersection pairing on $NS(S)$ is nondegenerate, e.g. if S is projective, then $T(S)$ is the orthogonal complement in $H^2(S, \mathbb{Z})$ of the Néron-Severi group.

Note that for a general K3 surface S we have no algebraic classes, i.e. $T(S) = H^2(S, \mathbb{Z})$. The *Picard number* $\rho(S)$ of S is the rank of the Néron-Severi group $NS(S)$.

For future use we also need the *Kähler cone* of a K3 surface, which is the cone of classes of Kähler $(1, 1)$ -forms. This lives in $H^{1,1}(S, \mathbb{R}) = H^{1,1}(S) \cap H^2(S, \mathbb{R})$. The restriction of the intersection product to $H^{1,1}(S, \mathbb{R})$ has signature $(1, 19)$. Let

$$\mathcal{C}_S \subset \{x \in H^{1,1}(S, \mathbb{R}) \mid (x, x) > 0\}$$

be the connected component that contains one (and hence all) Kähler classes. This is called the *positive cone* of S .

A class in $H^2(S, \mathbb{Z})$ is called *effective* if it is represented by an effective divisor. By a *nodal* class we mean the class δ of an effective divisor D of self-intersection $D^2 = -2$. We denote by Δ the set of all nodal classes. Every nodal class $\delta \in \Delta$ defines a reflection

$$s_\delta: H^2(S, \mathbb{Z}) \rightarrow H^2(S, \mathbb{Z})$$

given by $s_\delta(x) = x + (x, \delta)\delta$ and called the *Picard-Lefschetz reflection* defined by δ . We shall denote the \mathbb{R} - and \mathbb{C} -linear extensions of s_δ by the same symbol. Clearly s_δ is the reflection in the hyperplane H_δ orthogonal to δ . The set of effective classes on S is the semi-group generated by the integral points in the closure of \mathcal{C}_S and the nodal classes. The connected components of the set $\mathcal{C}_S \setminus \bigcup_{\delta \in \Delta} H_\delta$ are called the *chambers* of \mathcal{C}_S . The chamber

$$(2.2) \quad \mathcal{C}_S^+ = \{x \in \mathcal{C}_S \mid (x, \delta) > 0 \text{ for all effective } \delta \in \Delta\}$$

is equal to the Kähler cone of S according to [6, Corollary VIII.3.9].

2.2. Markings and the period map

A *marking* of a K3 surface is an isometry $\phi: H^2(S, \mathbb{Z}) \rightarrow L_{K3}$ and we refer to a pair (S, ϕ) as a *marked* K3 surface. An isomorphism between marked K3 surfaces is an isomorphism between the surfaces that commutes with the markings. If ω is a non-zero 2-form on S then $\mathbb{C}\phi(\omega) = \phi(H^{2,0}(S))$ is a line in the complex vector space $L_{K3} \otimes \mathbb{C}$.

For any indefinite lattice L we define

$$(2.3) \quad \Omega_L = \{[x] \in \mathbb{P}(L \otimes \mathbb{C}) \mid (x, x) = 0, (x, \bar{x}) > 0\}.$$

In the case of the K3 lattice, $\Omega = \Omega_{L_{K3}}$ is a connected complex manifold of dimension 20, called the *period domain of K3 surfaces*. Since $(\omega, \omega) = 0$ and $(\omega, \bar{\omega}) > 0$ it follows that $[\phi(\omega)] \in \Omega$. This is the *period point* of the marked K3 surface (S, ϕ) .

Let $p: \mathcal{S} \rightarrow \mathcal{U}$ be a flat family of K3 surfaces over some sufficiently small contractible open set \mathcal{U} . If $\phi_0: H^2(S_0, \mathbb{Z}) \rightarrow L_{K3}$ is a marking then this can be extended to a marking $\phi_{\mathcal{U}}: R^2p_*\mathbb{Z}_{\mathcal{U}} \rightarrow (L_{K3})_{\mathcal{U}}$ where $(L_{K3})_{\mathcal{U}}$ is the constant sheaf with fibre L_{K3} on \mathcal{U} . This defines a holomorphic map $\pi_{\mathcal{U}}: \mathcal{U} \rightarrow \Omega$, called the *period map* defined by the family $p: \mathcal{S} \rightarrow \mathcal{U}$.

2.3. The Torelli theorem

The Torelli problem asks how much information about an algebraic variety can be reconstructed from its Hodge structure. In the case of K3 surfaces this means whether one can recover a K3 surface S from a period point. This question can be made precise in different ways. As we shall see, one can prove a very strong form of the Torelli theorem in the case of K3 surfaces.

We start by discussing the local Torelli theorem. For this let $p: \mathcal{S} \rightarrow \mathcal{U}$ be a representative of the Kuranishi family (or versal deformation) of S . Since $H^0(S, T_S) = H^2(S, T_S) = 0$ the base space of the Kuranishi family is smooth of dimension $h^1(S, T_S) = h^{1,1}(S) = 20$. Choosing any marking of the central fibre of the Kuranishi family defines a marking for the entire family (we shall choose \mathcal{U} sufficiently small and contractible) and hence a period map $\pi: \mathcal{U} \rightarrow \Omega$.

Theorem 2.4. (Local Torelli) *The base space of the Kuranishi family is smooth of dimension 20. It is universal for all points in a sufficiently small neighbourhood of the origin. The differential of the period map is an isomorphism and thus the period map is a local isomorphism.*

Proof. See [6, Theorem VIII.7.3]. □

In order to discuss the global Torelli theorem we need the notion of Hodge isometry. Let S and S' be K3 surfaces. An isomorphism of \mathbb{Z} -modules $\Phi: H^2(S, \mathbb{Z}) \rightarrow H^2(S', \mathbb{Z})$ is called a *Hodge isometry* if it is an isometry and if its \mathbb{C} -linear extension $\Phi_{\mathbb{C}}: H^2(S, \mathbb{C}) \rightarrow H^2(S', \mathbb{C})$ preserves the Hodge decomposition. It is moreover called *effective* if it maps \mathcal{C}_S to \mathcal{C}'_S and maps effective classes to effective classes.

Proposition 2.5. *Let S and S' be K3 surfaces. Then the following are equivalent for a Hodge isometry $\Phi: H^2(S, \mathbb{Z}) \rightarrow H^2(S', \mathbb{Z})$:*

- (i) Φ is effective,
- (ii) $\Phi(\mathcal{C}_S^+) \subset \mathcal{C}'_S$, i.e. Φ maps the Kähler cone of S into that of S' .
- (iii) Φ maps one element of the Kähler cone of S into the Kähler cone of the surface S' .

Proof. See [6, Proposition VIII.3.11]. □

The crucial result for the theory of moduli of K3 surfaces is the following.

Theorem 2.6. (Strong Torelli) *Let S and S' be two K3 surfaces and suppose that $\Phi: H^2(S', \mathbb{Z}) \rightarrow H^2(S, \mathbb{Z})$ is an effective Hodge isometry. Then there is a unique isomorphism $f: S \rightarrow S'$ that induces Φ , i.e. such that $\Phi = f^*$.*

A proof of this theorem can be found in [6, Sections VIII.7–VIII.11]. Very roughly speaking the idea is to prove the Torelli theorem for projective Kummer surfaces first. The second step then consists of showing that the period points of marked Kummer surfaces are dense (in the complex topology) in the period domain Ω . The final step is then to prove the Torelli theorem for all K3 surfaces by approximating them by Kummer surfaces and taking suitable limits in the Barlet topology.

The following weaker form of the Torelli theorem is still useful.

Theorem 2.7. (Weak Torelli theorem) *Two K3 surfaces S and S' are isomorphic if and only if there is a Hodge isometry $H^2(S', \mathbb{Z}) \rightarrow H^2(S, \mathbb{Z})$.*

Proof. Assume that $\Phi: H^2(S', \mathbb{Z}) \rightarrow H^2(S, \mathbb{Z})$ is a Hodge isometry. Let W_S be the group of isometries of $H^2(S, \mathbb{Z})$ generated by the Picard-Lefschetz reflections. This group acts on the positive cone \mathcal{C}_S properly discontinuously. The closure of the Kähler cone in the positive cone is a fundamental domain for this action (cf. [6, Proposition VIII.3.10]). Hence we can choose an element $w \in W_S$ such that for a suitable choice of sign $\pm w \circ \Phi$ is an effective Hodge isometry by Proposition 2.5 and thus is induced by an isomorphism $f: S \rightarrow S'$ by Theorem 2.6. \square

2.4. The universal family of marked K3 surfaces

For each K3 surface S we choose a representative $p: \mathcal{S} \rightarrow \mathcal{U}$ of the Kuranishi family with \mathcal{U} contractible and sufficiently small such that the following hold:

- (i) $p: \mathcal{S} \rightarrow \mathcal{U}$ is the Kuranishi family for each point $s \in \mathcal{U}$.
- (ii) If $\phi: \mathbb{R}^2 p_* \mathbb{Z}_{\mathcal{S}} \rightarrow (L_{K3})_{\mathcal{U}}$ is a marking, then the associated period map $\pi: \mathcal{U} \rightarrow \Omega$ is injective.

We consider all marked Kuranishi families, i.e. pairs $(p: \mathcal{S} \rightarrow \mathcal{U}, \phi)$, having the above properties. We can glue the various copies of \mathcal{U} by identifying those points where the marked K3 surfaces are isomorphic. This defines a space M_1 all of whose points have neighbourhoods isomorphic to some \mathcal{U} . Hence M_1 is a 20-dimensional analytic manifold, but possibly not Hausdorff. It is also possible to show (cf. [6, Theorem VIII.10.6]) that one can glue the Kuranishi families to obtain a global family of K3 surfaces over M_1 .

It turns out that the space M_1 is indeed not Hausdorff. This follows from an example due to Atiyah, also referred to as Atiyah’s flop and nowadays crucial in higher-dimensional birational geometry. Consider, for example, the following 1-parameter family S_t of quartic (and hence K3) surfaces in \mathbb{P}^3 , which is given in affine coordinates by

$$x^2(x^2 - 2) + y^2(y^2 - 2) + z^2(z^2 - 2) = 2t^2.$$

Let the parameter t vary in a small neighbourhood B of the origin. For $t \neq 0$ these surfaces are smooth, whereas the surface S_0 has an ordinary double point at $(0,0,0)$. This is also the only singularity, again an ordinary double point, of the total space \mathcal{S} . Blowing up this node one obtains a smooth 3-fold $\tilde{\mathcal{S}}$ which contains a quadric $\mathbb{P}^1 \times \mathbb{P}^1$ as exceptional divisor E . The proper transform \hat{S}_0 of S_0 is a smooth K3 surface intersecting the exceptional divisor E in a rational curve of bidegree $(1, 1)$. This is a nodal curve on \hat{S}_0 . The two rulings on E can each be contracted giving rise to smooth 3-dimensional spaces $p_1: \mathcal{S}_1 \rightarrow B$ and $p_2: \mathcal{S}_2 \rightarrow B$. These families are by construction identical over $B \setminus 0$. They are, however, not identical over all of B , since the identity on $\mathcal{S} \setminus S_0$ would, otherwise, have to extend to an automorphism of the total space acting non-trivially on the tangent cone of the double point, which is clearly impossible. Now choose a marking for p_1 . This also defines a marking for p_2 . Since the families coincide outside the origin, the period maps π_1 and π_2 also

coincide away from 0. However, the markings differ for the central fibre (namely by the Picard-Lefschetz reflection defined by the nodal curve on \hat{S}_0). This shows that M_1 cannot be Hausdorff. In fact all non-Hausdorff behaviour comes from the existence of different resolutions of double points in families: see [20, Theorem 1' and Section 7].

There is another formulation of the Torelli theorem, which we will now describe. For this we consider

$$K\Omega = \{(\kappa, [\omega]) \in (L_{K3} \otimes \mathbb{R}) \times \Omega \mid (\kappa, \text{Re}(\omega)) = (\kappa, \text{Im}(\omega)) = 0, (\kappa, \kappa) > 0\}$$

and define $E(\kappa, \omega)$ as the oriented 3-dimensional space spanned by the ordered basis $\{\kappa, \text{Re}(\omega), \text{Im}(\omega)\}$. By mapping each point $(\kappa, [\omega])$ to the space $E(\kappa, \omega)$ we obtain a fibration

$$\Pi: K\Omega \rightarrow \text{Gr}^+(3, L_{K3} \otimes \mathbb{R})$$

over the Grassmannian $\text{Gr}^+(3, L_{K3} \otimes \mathbb{R})$ of oriented 3-planes in $L_{K3} \otimes \mathbb{R}$ on which the form $(\ , \)$ is positive definite. This is an $\text{SO}(3, \mathbb{R})$ -fibre bundle and the projection Π is equivariant with respect to the action of the orthogonal group $\text{Aut}(L_{K3} \otimes \mathbb{R}) \cong \text{O}(3, 19)$. We define

$$(K\Omega)^0 := \{(\kappa, [\omega]) \in K\Omega \mid (\kappa, d) \neq 0 \text{ for } d \in L_{K3}, (d, d) = -2, (\omega, d) = 0\}.$$

This is an open subset of $K\Omega$.

For every point $\omega \in \Omega$ we consider the cone

$$C_\omega = \{x \in L_{K3} \otimes \mathbb{R} \mid (x, \omega) = 0, (x, x) > 0\}.$$

This has two connected components and varies differentiably with ω . Since Ω is connected and simply connected, we can globally choose one of these components, say C_ω^+ . Let (S, ϕ) be a marked K3 surface and let $\kappa \in H^{1,1}(S, \mathbb{R})$ be a Kähler class. Then we say that (S, κ) , or more precisely $((S, \phi), \kappa)$, is a *marked pair* if $\phi_C(\kappa) \in C_\omega^+$, where ω is the period point defined by (S, ϕ) .

Let M_2' be the real analytic vector bundle with fibre $H^{1,1}(S_t)$ over the base M_1 of the universal family of marked K3 surfaces and let $M_2 \subset M_2'$ be the subset of Kähler classes. This is open by [72, Theorem 15]. In particular, M_2 is real analytic of dimension $60 = 40 + 20$, where 40 is the real dimension of the base M_1 and 20 is the dimension of the fibre. We can now define a real-analytic map

$$\pi_2: M_2 \rightarrow (K\Omega)^0$$

by mapping $\kappa \in H^{1,1}(S_t), t \in M_1$ to $\pi_2(\kappa) = (\phi_C(\kappa), \pi(t))$. This is called the *refined period map*. In this way we obtain the obvious commutative diagram

$$\begin{array}{ccc} M_2 & \xrightarrow{\pi_2} & (K\Omega)^0 \\ \downarrow & & \downarrow \\ M_1 & \xrightarrow{\pi_1} & \Omega. \end{array}$$

The Torelli theorem can now be reformulated as follows.

Theorem 2.8. *The map π_2 is injective (and thus M_2 is Hausdorff).*

Using this one can finally prove that the period map is surjective.

Theorem 2.9. (Surjectivity of the period map) *The refined period map π_2 is surjective. In particular, every point of Ω appears as the period point of some marked K3 surface.*

Proof. This is proven in [6, Section VIII.14]. □

The surjectivity of the period map was one of the major questions in the theory of K3 surfaces. A. Todorov [126] was the first to give a proof; the argument given in [6] is due to Looijenga.

In view of Theorem 2.7 and Theorem 2.9 we now have

Theorem 2.10. *The set $O(L_{K3}) \backslash \Omega$ is in 1 : 1 correspondence with the set of isomorphism classes of K3 surfaces.*

We can thus think of $O(L_{K3}) \backslash \Omega$ as the “moduli space of K3 surfaces”. It must be pointed out, however, that the action of the group $O(L_{K3})$ is not well behaved. In particular, it is not properly discontinuous. We shall now turn to the case of polarised K3 surfaces where we shall see that the situation is much better.

2.5. Moduli spaces of polarised K3 surfaces

A *polarisation* on a K3 surface is an ample line bundle \mathcal{L} . Since the irregularity of K3 surfaces is 0 and the Picard group has no torsion we can identify a line bundle \mathcal{L} with its first Chern class $h = c_1(\mathcal{L}) \in H^2(S, \mathbb{Z})$. An ample line bundle \mathcal{L} is nef and big, and conversely a nef and big line bundle on a K3 surface is ample if there are no (-2) -curves C with $h \cdot C = 0$. This can be seen from the description of the Kähler cone (2.2) and Nakai’s criterion, or from Reider’s theorem [6, Theorem IV.11.4] or from [113]. Throughout, we shall only consider primitive polarisations, i.e. we shall assume that the class h is non-divisible in the K3 lattice. The *degree* of a polarisation is $\text{deg}(\mathcal{L}) = h^2 = 2d$. Note that the degree is always even.

We denote by $\langle -2d \rangle$ the rank 1 lattice whose generator has square $-2d$.

Lemma 2.11. *Suppose $h \in L_{K3}$ is a primitive vector with $h^2 = 2d > 0$. Then the orthogonal complement $L_h = h^\perp_{L_{K3}}$ of h is isometric to L_{2d} , where L_{2d} , the lattice L_{2d} associated with K3 surfaces with a polarisation of degree $2d$, is defined by*

$$(2.12) \quad L_{2d} = 2U \oplus 2E_8(-1) \oplus \langle -2d \rangle.$$

Proof. It follows from Eichler’s criterion (see Lemma 7.8 and Example 7.9 below) that there is a unique $O(L_{K3})$ -orbit of primitive vectors h of given degree in the K3 lattice $L_{K3} = 3U \oplus 2E_8(-1)$. Hence we can assume that h is in one of the copies of the hyperbolic plane and that $h = e + df$ where e, f are a basis of a copy of U with $e^2 = f^2 = 0$ and $(e, f) = 1$. The structure of L_h is clear in this case. □

The lattice L_{2d} is an even lattice of signature $(2, 19)$. The period domain $\Omega_{2d} = \Omega_{L_{2d}}$ has two connected components, \mathcal{D}_{2d} and \mathcal{D}'_{2d} , interchanged by complex conjugation. The domain \mathcal{D}_{2d} is a 19-dimensional symmetric homogeneous domain of type IV: see [114, Appendix 6]. One can also describe Ω_{2d} as the intersection of the domain Ω with the hyperplane $h_{L_{K3}}^\perp$.

We shall fix $h \in L_{K3}$ once and for all. For each polarised K3 surface (S, \mathcal{L}) of degree $2d$ we can consider polarised markings, i.e. markings $\phi: H^2(S, \mathbb{Z}) \rightarrow L_{K3}$ with $\phi(c_1(\mathcal{L})) = h$. Any two such markings differ by an element in the group

$$(2.13) \quad O(L_{K3}, h) = \{g \in O(L_{K3}) \mid g(h) = h\}.$$

This group leaves the orthogonal complement L_{2d} of h invariant and hence is a subgroup of $O(L_{2d})$.

For any lattice L we denote by $L^\vee = \text{Hom}(L, \mathbb{Z})$ its dual lattice. The *discriminant group* $D(L) = L^\vee/L$ is a finite abelian group of order $|\det L|$ and carries a discriminant quadratic form q_L (if L is even) and a discriminant bilinear form b_L , with values in $\mathbb{Q}/2\mathbb{Z}$ and \mathbb{Q}/\mathbb{Z} respectively (see [97, Section 1.3]). The *stable orthogonal group* of an even lattice L is defined as the kernel

$$(2.14) \quad \tilde{O}(L) = \ker(O(L) \rightarrow O(D(L))).$$

If L is indefinite, the subgroup $O^+(L)$ is defined to be the group of elements of real spinor norm 1 (see [49] for a definition of the real spinor norm). We define

$$(2.15) \quad \tilde{O}^+(L) = \tilde{O}(L) \cap O^+(L)$$

and generally, for any subgroup $G < O(L)$, we use G^+ to denote the kernel of the real spinor norm and \tilde{G} to denote the stable subgroup, the kernel of $G \rightarrow O(D(L))$.

For $h \in L_{K3}$ with $h^2 = 2d$, it follows from Nikulin’s theory [97, Corollary 1.5.2] that

$$O(L_{K3}, h) = \tilde{O}(L_{2d})$$

considered as subgroups of $O(L_{2d})$. The two connected components \mathcal{D}_{2d} and \mathcal{D}'_{2d} are interchanged by the group $O(L_{2d})$. The index 2 subgroup that fixes the components is $O^+(L_{2d})$. Finally we define

$$(2.16) \quad \mathcal{F}_{2d} = \tilde{O}(L_{2d}) \backslash \Omega_{2d} = \tilde{O}^+(L_{2d}) \backslash \mathcal{D}_{2d}.$$

It is important to note that the situation is much better here than in the non-polarised case. The reason lies in the change of signature which is now $(2, 19)$ rather than $(3, 19)$. We are thus dealing with locally symmetric hermitian domains and, as a result, the action of the group $O(L_{2d})$ on Ω_{2d} is properly discontinuous. Hence the quotient space of Ω_{2d} by any subgroup of $O(L_{2d})$ of finite index (i.e. an *arithmetic subgroup*) is a complex space with only finite quotient singularities (also sometimes called a V-manifold). By a famous result of Baily and Borel [5] these quotients are quasi-projective and thus carry a natural structure of an algebraic variety. We

shall discuss various compactifications of these quasi-projective varieties below in Section 5.

In order to describe the moduli space of polarised K3 surfaces we need one more ingredient. For $h \in L_{K3}$ we define

$$\Delta_h = \{\delta \in L_{K3} \mid \delta^2 = -2, (\delta, h) = 0\}.$$

For each $\delta \in \Delta_h$ we define the hyperplane $H_\delta = \delta^\perp_{L_{K3}}$, i.e. the hyperplane fixed by the Picard-Lefschetz reflection defined by δ . We set

$$\Omega_{2d}^0 = \Omega_{2d} \setminus \bigcup_{\delta \in \Delta_h} (H_\delta \cap \Omega_{2d}).$$

There are only finitely many $\tilde{O}(L_{2d})$ -orbits in Δ_h . This follows immediately from Lemma 7.8, below: in fact there are at most two orbits by [48, Proposition 2.4(ii)]. Since the group acts properly discontinuously on Ω , the hyperplanes H_δ for $\delta \in \Delta_h$ form a locally finite collection. Clearly, the action of the group $\tilde{O}(L_{2d})$ on Ω_{2d} restricts to an action on Ω_{2d}^0 . We define

$$\mathcal{F}_{2d}^0 = \tilde{O}(L_{2d}) \backslash \Omega_{2d}^0.$$

Note that this is again a quasi-projective variety (it arises from \mathcal{F}_{2d} by removing finitely many divisors) with only finite quotient singularities.

Theorem 2.17. *The variety \mathcal{F}_{2d}^0 is the moduli space of polarised K3 surfaces of degree $2d$, i.e. its points are in $1 : 1$ correspondence with polarised K3 surfaces of degree $2d$.*

Proof. Let (S, \mathcal{L}) be a polarised K3 surface with $\deg(\mathcal{L}) = 2d$. We consider polarised markings $\phi: H^2(S, \mathbb{Z}) \rightarrow L_{K3}$ with $\phi(c_1(\mathcal{L})) = h$. Since $(\omega_S, c_1(\mathcal{L})) = 0$ we find that $\phi(\omega_S) \in \Omega_{2d}$. In fact, since \mathcal{L} is ample it has positive degree on all (-2) -curves and hence $\phi(\omega_S)$ lies in Ω_{2d}^0 . Any two polarised markings differ by an element in $O(L_{K3}, h) = \tilde{O}(L_{2d})$ and hence we obtain a well-defined map which associates to an isomorphism class (S, \mathcal{L}) a point in \mathcal{F}_{2d}^0 . This map is injective: assume that a point in \mathcal{F}_{2d}^0 arises from two polarised surfaces (S, \mathcal{L}) and (S', \mathcal{L}') . Then there exists a Hodge isometry $H^2(S', \mathbb{Z}) \rightarrow H^2(S, \mathbb{Z})$ mapping the ample class $c_1(\mathcal{L}')$ to $c_1(\mathcal{L})$. It then follows from the strong Torelli Theorem (Theorem 2.6) that this map is induced by an isomorphism $(S, \mathcal{L}) \cong (S', \mathcal{L}')$. Thus we get an injective map from the set of isomorphism classes of degree $2d$ polarised K3 surfaces into \mathcal{F}_{2d}^0 . Finally, the surjectivity of this map follows from the surjectivity of the period map, Theorem 2.9. \square

In the literature one often finds references to \mathcal{F}_{2d} as the moduli space of polarised K3 surfaces. One can interpret the points in the complement of \mathcal{F}_{2d}^0 as weakly- or semi-polarised K3 surfaces, i.e. \mathcal{L} has positive degree and is nef, but not ample, as it has degree 0 on some nodal class(es). Alternatively, one can consider ample line bundles on K3 surfaces with rational double points. There is still a version of the strong Torelli theorem for weakly polarised K3 surfaces due to D. Morrison [88],

but the precise formulation is subtle. For purposes of the birational geometry of these spaces, it obviously does not matter whether one works with \mathcal{F}_{2d} or its open part \mathcal{F}_{2d}^0 .

The notion of polarised K3 surfaces was generalised by Nikulin [96] to that of *lattice-polarised* K3 surfaces: see also Dolgachev’s paper [26] for a concise account, in particular in connection with mirror symmetry. To describe this, we fix a lattice M of signature $(1, t)$, which we assume can be embedded primitively into the K3-lattice L_{K3} . The cone $V_M = \{x \in M_{\mathbb{R}} \mid (x, x) > 0\}$ has two connected components: we fix one and denote it by \mathcal{C}_M . Let

$$\Delta_M = \{d \in M \mid (d, d) = -2\}$$

and choose a decomposition $\Delta_M = \Delta_M^+ \cup (-\Delta_M^+)$. Moreover let

$$\mathcal{C}_M^+ = \{h \in V(M)^+ \cap M \mid (h, d) > 0 \text{ for all } d \in \Delta_M^+\}.$$

An *M-polarised* K3 surface is then a pair (S, j) where S is a K3 surface and $j: M \hookrightarrow \text{Pic}(S)$ is a primitive embedding. An isomorphism between M -polarised K3 surfaces is an isomorphism between the surfaces that commutes with the primitive embeddings. We call (S, j) *ample* (or *pseudo-ample*), if $j(\mathcal{C}_M^+)$ contains an ample (or pseudo-ample) class. The classical case of polarised K3 surfaces is the case where $t = 0$ and $M = \langle 2d \rangle$.

The theory of moduli of polarised K3 surfaces carries over naturally to lattice polarised K3 surfaces. For this, one has to consider the domain $\Omega_M = \{\omega \in \Omega \mid (w, M) = 0\}$ and the group $O(L_{K3}, M)$ of orthogonal transformations of the K3 lattice which fixes M . The role of the variety \mathcal{F}_{2d} is then taken by the quotient

$$(2.18) \quad \mathcal{F}_M = O(L_{K3}, M) \backslash \Omega_M.$$

Lattice polarised K3 surfaces play a role in mirror symmetry. For this we consider *admissible* lattices, i.e. lattices M which admit an embedding $j: M \hookrightarrow L_{K3}$ such that

$$M_{L_{K3}}^{\perp} \cong \mathfrak{u}(\mathfrak{m}) \oplus \widetilde{M}.$$

The choice of such a splitting determines a primitive embedding $\tilde{j}: \widetilde{M} \hookrightarrow L_{K3}$. The lattice \widetilde{M} is hyperbolic: more precisely, its signature is $(1, 18 - t)$. The variety $\mathcal{F}_{\widetilde{M}}$ is then a mirror family to \mathcal{F}_M . There are more ingredients to the concept of mirror symmetry which we will not describe here, such as the Yukawa coupling and the mirror map. For details we refer to [26]: see also [52], [56].

Finally, we want to comment on the relationship between the construction of moduli spaces of K3 surfaces as quotients of homogeneous domains and GIT-like constructions. By Viehweg’s results [128] moduli spaces of polarised varieties with trivial canonical bundle exist and can be constructed as group quotients. For this we first fix a Hilbert polynomial $P(m)$ of a line bundle on a K3 surface. This is of the form $P(m) = m^2 d + 2$ where the degree of the line bundle is $2d$. Let \mathcal{M}_{2d} be the GIT moduli space of degree $2d$ polarised K3 surfaces. We want to relate this to \mathcal{F}_{2d} .

We first note that for any ample line bundle \mathcal{L} on a K3 surface S its third power $\mathcal{L}^{\otimes 3}$ is very ample. For this, see [113], but in general one can use Matsusaka's big theorem ([87], [79]) to show that there is a positive integer m_0 such that $\mathcal{L}^{\otimes m_0}$ is very ample for all polarised varieties (X, \mathcal{L}) with fixed Hilbert polynomial. Now let $m_0 \geq 3$ be sufficiently big. Then we have embeddings $f_{|\mathcal{L}^{\otimes m_0}|}: S \rightarrow \mathbb{P}^{N-1}$ where $N = h^0(S, \mathcal{L}^{\otimes m_0}) = P(m_0)$. Such an embedding depends on the choice of a basis of $H^0(S, \mathcal{L}^{\otimes m_0})$. Let H be an irreducible component of the Hilbert scheme $\text{Hilb}_{\mathbb{P}}(\mathbb{P}^{N-1})$ containing at least one point corresponding to a smooth K3 surface S . Let H_{sm} be the open part of H parametrising smooth surfaces. Then it is easy to prove that H_{sm} is smooth and that every point in H_{sm} parametrises a K3 surface. There exists a universal family $\mathcal{S}_{\text{sm}} \rightarrow H_{\text{sm}}$. The group $SL(N, \mathbb{C})$ acts on H_{sm} and every irreducible component of the GIT moduli space of degree $2d$ polarised K3 surfaces is of the form $SL(N, \mathbb{C}) \backslash H_{\text{sm}}$. Let \mathcal{M}'_{2d} be such a component. Choosing local polarised markings for the universal family one can construct a map $H_{\text{sm}} \rightarrow \mathcal{F}_{2d}^0$, which clearly factors through the action of $SL(N, \mathbb{C})$, i.e. gives rise to a map $\pi: \mathcal{M}'_{2d} \rightarrow \mathcal{F}_{2d}^0$. By construction this is a holomorphic map. On the other hand both \mathcal{M}'_{2d} and \mathcal{F}_{2d}^0 are quasi-projective varieties. It then follows from a theorem of Borel [15] that π is a morphism of quasi-projective varieties.

We claim that π is an isomorphism and that \mathcal{M}_{2d} has only one component. First of all we note that one can, as in the proof of [50, Theorem 1.5], take a finite étale cover $H'_{\text{sm}} \rightarrow H_{\text{sm}}$ such that the action of $SL(N+1, \mathbb{C})$ lifts to a free action on H'_{sm} as well as on the pullback $\mathcal{S}'_{\text{sm}} \rightarrow H'_{\text{sm}}$ of the universal family. This gives a quotient family over $Z_{\text{sm}} = SL(N+1, \mathbb{Z}) \backslash H'_{\text{sm}}$ which is smooth and maps finite-to-one to \mathcal{M}_{2d} . By the local Torelli theorem the natural map $Z_{\text{sm}} \rightarrow \mathcal{F}_{2d}^0$ has discrete fibres and hence the same is true for π . But then π must be dominant. Now we can use Theorem 2.17 to conclude that \mathcal{M}_{2d} is irreducible and that π is a bijection. Since \mathcal{F}_{2d}^0 is a normal variety, it also follows that π is an isomorphism. We can thus summarise our discussion as follows.

Theorem 2.19. *There is an isomorphism $\mathcal{M}_{2d} \cong \mathcal{F}_{2d}^0$, i.e. the GIT moduli space \mathcal{M}_{2d} is isomorphic to the modular variety \mathcal{F}_{2d}^0 .*

3. Irreducible symplectic manifolds

In this section we recall the main properties of irreducible symplectic manifolds, discuss the Torelli theorem and give basic facts about moduli spaces of polarised symplectic manifolds.

3.1. Basic theory of irreducible symplectic manifolds

The theory of irreducible symplectic manifolds is less developed than that of K3 surfaces. Nevertheless, several results have been proved over the last 30 years.

Definition 3.1. A complex manifold X is called an *irreducible symplectic manifold* or *hyperkähler manifold* if the following conditions are fulfilled:

- (i) X is a compact Kähler manifold;
- (ii) X is simply-connected;
- (iii) $H^0(X, \Omega_X^2) \cong \mathbb{C}\omega$ where ω is an everywhere nondegenerate holomorphic 2-form.

According to the Bogomolov decomposition theorem [9], irreducible symplectic manifolds are one of the building blocks for compact Kähler manifolds with trivial canonical bundle: see also [8, Théorème 2]. The others are abelian varieties and Calabi-Yau varieties (here we mean Calabi-Yau in its strictest sense, i.e. a compact Kähler manifold X such that $\pi_1(X) = 1$ and $H^0(X, \Omega_X^i) = 0$ for $0 < i < \dim X$).

In dimension two the only irreducible symplectic manifolds are K3 surfaces. Although irreducible symplectic manifolds have now been studied for nearly 30 years, only four classes of such manifolds have so far been discovered and it is a wide open problem whether other types exist or not. The known examples are:

(i) The length n Hilbert scheme $S^{[n]} = \text{Hilb}^n(S)$ for a K3 surface S , and its deformations. Note that the deformation space of such a variety has dimension 21 if $n \geq 2$ and that, since K3 surfaces only depend on 20 parameters, a general deformation will not itself be of the form $S^{[n]}$. We shall refer to these varieties as irreducible symplectic manifolds of *deformation K3^[n] type* or *deformation K3^[n] manifolds*.

(ii) Let A be a 2-dimensional complex torus and consider the length- $(n + 1)$ Hilbert scheme $A^{[n+1]} = \text{Hilb}^{n+1}(A)$ together with the morphism $p: A^{[n+1]} \rightarrow A$ given by addition. Then $X = p^{-1}(0)$ is an irreducible symplectic manifold, called a *generalised Kummer variety* (even though it is not necessarily algebraic). The deformation space of these manifolds has dimension 5 if $n \geq 2$, again one more than for 2-dimensional complex tori.

(iii) O’Grady’s irreducible symplectic manifolds of dimension 6, described in [100]. These are deformations of (desingularised) moduli spaces of sheaves on an abelian surface and depend on 6 parameters.

(iv) O’Grady’s irreducible symplectic manifolds of dimension 10, described in [99]. These are deformations of (desingularised) moduli spaces of sheaves on a K3 surface and depend on 22 parameters.

Other moduli spaces of sheaves, apart from those considered in [99] and [100], cannot be desingularised symplectically: see [67] and also [132].

In many ways irreducible symplectic manifolds behave like K3 surfaces, but there are also important differences, as we shall see later. We first notice that it follows immediately from the definition that X must have even dimension $2n$ over \mathbb{C} and that its canonical bundle ω_X is trivial (an n -fold exterior power of a generator ω of $H^0(X, \Omega_X^2)$ will define a trivialisation of the canonical bundle). Clearly $h^{2,0}(X) = h^{0,2}(X) = 1$ and $h^{1,0}(X) = h^{0,1}(X) = 0$. By a result of Bogomolov [10],

the deformation space of X is unobstructed. This result was generalised to Ricci-flat manifolds by Tian [125] and Todorov [126], and algebraic proofs were given by Kawamata [68] and Ran [109] (see also [36]). Since

$$T_{[0]} \text{Def}(X) \cong H^1(X, T_X) \cong H^1(X, \Omega_X^1)$$

the dimension of the deformation space is $b_2(X) - 2$.

As in the K3 case we have a Hodge decomposition $H^2(X, \mathbb{C}) = H^{2,0} \oplus H^{1,1} \oplus H^{0,2}$ with $H^{2,0}$ and $H^{0,2}$ both 1-dimensional. Unlike the K3 case the intersection form does not immediately provide $H^2(S, \mathbb{Z})$ with the structure of a lattice. It was, however, shown by Beauville [8] that $H^2(X, \mathbb{Z})$ does carry a natural structure as a lattice. To define this, let $\omega \in H^{2,0}(X)$ be such that $\int_X (\omega \bar{\omega})^n = 1$ and define

$$q'_X(\alpha) = \frac{n}{2} \int_X \alpha^2 (\omega \bar{\omega})^{n-1} + (1-n) \left(\int_X \alpha \omega^{n-1} \bar{\omega}^n \right) \left(\int_X \bar{\alpha} \omega^n \bar{\omega}^{n-1} \right).$$

After multiplication by a positive constant γ the quadratic form $q_X = \gamma q'_X$ defines an indivisible integral symmetric bilinear form $(\ , \)_X$ on $H^2(X, \mathbb{Z})$: this is the Beauville form. Clearly $(\omega, \omega)_X = 0$ and $(\omega, \bar{\omega})_X > 0$.

There is another way to introduce the Beauville form. For this let $v(\alpha) = \alpha^{2n}$ be given by the cup product. Then, by a result of Fujiki [36, Theorem 4.7], there is a positive rational number c , the *Fujiki invariant*, such that

$$v(\alpha) = c q_X(\alpha)^n$$

for all $\alpha \in H^2(X, \mathbb{Z})$. In this sense the Beauville form can be derived from the cup product of the cohomology.

Proposition 3.2. *The Beauville lattices and Fujiki invariants for the known examples of irreducible symplectic manifolds are as follows:*

(i) *The Beauville lattice of a deformation $K3^{[n]}$ manifold is $L_{K3, 2n-2} = 3\mathbb{U} \oplus 2E_8(-1) \oplus \langle -2(n-1) \rangle$. It has rank 23, one more than the K3 lattice, to which it is closely related. If $X = S^{[n]}$ for a K3 surface S , then $3\mathbb{U} \oplus 2E_8(-1)$ comes from $H^2(S, \mathbb{Z})$ and the summand $\langle -2(n-1) \rangle$ is generated (over \mathbb{Q}) by the exceptional divisor E , which is the blow-up of the diagonal in the symmetric product $S^{(n)}$. As an element in the Picard group the divisor E is 2-divisible. The Beauville lattice remains constant under deformations. The Fujiki invariant is $c = (2n)!/(n!2^n)$.*

(ii) *The Beauville lattice of a generalised Kummer variety (or deformation thereof) is $3\mathbb{U} \oplus \langle -2(n+1) \rangle$ and the Fujiki invariant is $c = (n+1)(2n)!/(n!2^n)$.*

(iii) *The Beauville lattice of the 6-dimensional example of O’Grady is $3\mathbb{U} \oplus \langle -2 \rangle \oplus \langle -2 \rangle$ and the Fujiki invariant is $c = 60$.*

(iv) *The Beauville lattice of the 10-dimensional example of O’Grady is $3\mathbb{U} \oplus 2E_8(-1) \oplus A_2(-1)$ and the Fujiki invariant is $c = 945$.*

For proofs, see [8], [111] and [110]. Note that all these lattices are even. It is, however, not known whether this is a general fact for irreducible symplectic manifolds.

Fix an abstract lattice L that is isomorphic to $(H^2(X, \mathbb{Z}), q_X)$ for some irreducible symplectic manifold X . A *marking* is an isomorphism of lattices $\phi: H^2(X, \mathbb{Z}) \xrightarrow{\sim} L$. Let $p: \mathcal{X} \rightarrow \mathcal{U}$ be a representative of the Kuranishi family of deformations of X with sufficiently small and contractible base. Note that by unobstructedness the base space of the Kuranishi family is smooth and of dimension $b_2(X) - 2$. The marking ϕ for X defines a marking for \mathcal{X} and we obtain a period map $\pi_{\mathcal{U}}: \mathcal{U} \rightarrow \Omega_L$ to the period domain defined by (2.3). As in the K3 case we have a local Torelli theorem.

Theorem 3.3. (Beauville) *The differential of the period map defined by the Kuranishi family is an isomorphism and thus the period map is a local isomorphism.*

Proof. See [9], [8]. □

As in the K3 case one can define a moduli space of marked irreducible symplectic manifolds (of a given type). Again, this will not be Hausdorff. Another result which carries over from the K3 case is surjectivity of the period map.

Theorem 3.4. (Huybrechts) *Let L be a lattice of an irreducible symplectic manifold and let Ω_L be the associated period domain. If \mathcal{M}'_L is a non-empty component of the moduli space \mathcal{M}_L of irreducible symplectic manifolds with Beauville lattice L , then the period map $\pi: \mathcal{M}'_L \rightarrow \Omega_L$ is surjective.*

Proof. A proof can be found in [63, Section 8]. □

3.2. Hodge theoretic Torelli theorem

So far, many results from K3 surfaces have carried over to other irreducible symplectic manifolds. The situation changes when it comes to the global Torelli theorem. The first counterexample to this is due to Debarre [24]. He showed the following: let S be a K3 surface containing only one curve, which is a smooth rational curve C , and consider the Hilbert scheme $X = S^{[n]}$ with $n \geq 2$. Then X contains $S^n(C) \cong \mathbb{P}^n$. One can perform an elementary transformation on X by first blowing up $S^n(C)$ and then contracting the exceptional divisor in another direction. This gives another compact complex manifold X' which is bimeromorphic but not isomorphic to X . In general X' need not be Kähler, but Debarre produced an example where X' does have a Kähler structure and thus is an irreducible symplectic manifold.

Since the natural bimeromorphic transformation $f: X' \dashrightarrow X$ defines a Hodge isometry $f^*: H^2(X, \mathbb{Z}) \rightarrow H^2(X', \mathbb{Z})$ this gives a counterexample to the global Torelli theorem. It should be noted, though, that neither the surface S nor the varieties X and X' are projective. Moreover, the existence of (-2) -curves on a K3 surface S is exactly the cause for the failure of the Hausdorff property for the base of the universal family.

Debarre's counterexample would still allow for a version of the Torelli theorem where the existence of a Hodge isometry only implies birational equivalence, not an isomorphism (for K3 surfaces the two notions coincide). However, this is also ruled

out by the following counterexample which is due to Y. Namikawa [95]. For this one starts with a generic abelian surface A with a polarisation of type $(1,3)$. Then the dual abelian surface \widehat{A} also carries a $(1,3)$ -polarisation. Let $X = \text{Km}^{[2]} A$ and $\widehat{X} = \text{Km}^{[2]}(\widehat{A})$ be the associated generalised Kummer varieties of dimension 4. Then X and \widehat{X} are birationally equivalent if and only if A and \widehat{A} are isomorphic abelian surfaces. The reason for this is the following: every birational isomorphism must send the exceptional divisor E on X to the exceptional divisor \widehat{E} on \widehat{X} . Since the Albanese of E and \widehat{E} are A and \widehat{A} respectively, this implies that A and \widehat{A} are isomorphic. This is not the case for general A . This shows that Namikawa's example gives a counterexample even to the birational global Torelli theorem. Moreover, one can easily make this into a counterexample to the polarised Torelli theorem. This can be done by choosing polarisations of the form $mL - \delta$ and $m\widehat{L} - \widehat{\delta}$, where m is sufficiently large, L and \widehat{L} are induced from the polarisation on A and \widehat{A} respectively, $\delta = 2E$ and $\widehat{\delta} = 2\widehat{E}$ (the exceptional divisors are 2-divisible in the Picard group). The Hodge isomorphism respects these polarisations.

At first, these counterexamples seem to indicate that there is no chance of proving a version of the global Torelli theorem for irreducible symplectic manifolds. However, the above example is not as surprising as it seems at a first glance. It is well known, and also well understood in terms of period domains and arithmetic groups, that A and \widehat{A} are not isomorphic as polarised abelian surfaces (their period points in the Siegel space are inequivalent under the paramodular group), but that the associated Kummer surfaces $\text{Km}(A)$ and $\text{Km}(\widehat{A})$ are isomorphic (and their period points in the corresponding type IV domain are equivalent under the orthogonal group). Details can be found in [43]. An analysis of this situation suggests that a version of the Torelli theorem could hold if one considers Hodge isometries with extra conditions.

Verbitsky [127] has announced a global Torelli theorem for irreducible symplectic manifolds (see also Huybrecht's Bourbaki talk [64]). His results were further elucidated by Markman [86]. The crucial idea here is to use monodromy operators and parallel transport operators. Markman first noticed the importance of these operators for the study of irreducible symplectic manifolds, developing his ideas in a series of papers [84], [83], [85]. To define them, let X_1, X_2 be irreducible symplectic manifolds. We say that $f: H^*(X_1, \mathbb{Z}) \rightarrow H^*(X_2, \mathbb{Z})$ is a *parallel transport operator* if there exists a smooth, proper flat family $\pi: \mathcal{X} \rightarrow B$ of irreducible symplectic manifolds together with points $b_1, b_2 \in B$ such that there are isomorphisms $\alpha_i: X_i \xrightarrow{\sim} \mathcal{X}_{b_i}$ and a continuous path $\gamma: [0,1] \rightarrow B$ with $\gamma(0) = b_1$ and $\gamma(1) = b_2$, such that the parallel transport in the local system $R\pi_*\mathbb{Z}$ along γ induces the isomorphism $(\alpha_2^{-1})^* \circ f \circ \alpha_1^*: H^*(\mathcal{X}_{b_1}, \mathbb{Z}) \rightarrow H^*(\mathcal{X}_{b_2}, \mathbb{Z})$.

For a single irreducible symplectic manifold X , we call an automorphism $f: H^*(X, \mathbb{Z}) \rightarrow H^*(X, \mathbb{Z})$ a *monodromy operator* if it is a parallel transport operator (with $X_1 = X_2 = X$). The *monodromy group* $\text{Mon}(X)$ is defined as the subgroup

of $GL(H^*(X, \mathbb{Z}))$ generated by monodromy operators. Restricting the group action to the second cohomology group we obtain a subgroup $\text{Mon}^2(X)$ of $GL(H^2(X, \mathbb{Z}))$. Since monodromy operators preserve the Beauville form we obtain a subgroup $\text{Mon}^2(X) \subset O(H^2(X, \mathbb{Z}))$.

Based on Verbitsky’s results [127], Markman [86] has formulated the following Hodge theoretic global Torelli theorem.

Theorem 3.5. (Hodge theoretic Torelli) *Suppose that X and Y are irreducible symplectic manifolds.*

- (i) *If $f: H^2(Y, \mathbb{Z}) \rightarrow H^2(X, \mathbb{Z})$ is an isomorphism of integral Hodge structures which is a parallel transport operator, then X and Y are bimeromorphic.*
- (ii) *If, moreover, f maps a Kähler class of Y to a Kähler class of X , then X and Y are isomorphic.*

Proof. The first part of the theorem follows easily from Verbitsky’s results. The second part uses in addition results on the Kähler cone of irreducible symplectic manifolds. For a more detailed discussion see [86, Theorem 1.3] and [86, Section 3.2]. □

3.3. Moduli spaces of polarised irreducible symplectic manifolds

We shall now turn to the case of polarised irreducible symplectic manifolds. In the K3 case we saw that the degree is the only invariant of a polarisation, or, equivalently, that there is only one $O(L_{K3})$ -orbit of primitive vectors of given length. This is no longer true in general, as can already be seen in the case of $S^{[2]}$. Recall that the Beauville lattice in this case is isomorphic to $L_{K3,2} = 3U \oplus 2E_8(-1) \oplus \langle -2 \rangle$. If h is a primitive vector the number $\text{div}(h)$ (the divisor of h : see Equation (7.4) below) is the positive generator of the ideal $(h, L_{K3,2})$, which is the biggest positive integer by which one can divide h as a vector in the dual lattice $L_{K3,2}^\vee$. Since $L_{K3,2}$ is not unimodular, but has determinant 2, the divisor $\text{div}(h)$ can be 1 or 2. Indeed, both of these happen when $d \equiv -1 \pmod{4}$ and accordingly we have one $O(L_{K3,2})$ -orbit if $d \not\equiv -1 \pmod{4}$ and two if $d \equiv -1 \pmod{4}$. Details of this can be found in [50]. These two cases are referred to as the *split* case ($\text{div}(h) = 1$) and the *non-split* case ($\text{div}(h) = 2$). The reason for this terminology is the behaviour of the orthogonal lattice: if $h^2 = 2d$ the possibilities for $L_h = h_{L_{K3,2}}^\perp$ are (see Example 7.12 below)

$$(3.6) \quad L_h = 2U \oplus 2E_8(-1) \oplus \langle -2 \rangle \oplus \langle -2d \rangle \text{ for } \text{div}(h) = 1$$

and

$$(3.7) \quad L_h = 2U \oplus 2E_8(-1) \oplus \begin{pmatrix} -2 & 1 \\ 1 & -\frac{d+1}{2} \end{pmatrix} \text{ for } \text{div}(h) = 2.$$

For the higher dimensional case of $S^{[n]}$ the situation becomes more involved as the possibilities for the divisor of h increase in number. Whenever the primitive vectors of length d form more than one orbit, the moduli space of polarised irreducible symplectic manifolds of degree d will not be connected: see [50].

In order to discuss moduli spaces of polarised irreducible symplectic manifolds we first fix some discrete data: the dimension $2n$, the Beauville lattice (considered as an abstract lattice L), and the Fujiki invariant c . Together L and c define the *numerical type* of the irreducible symplectic manifold, denoted by \mathbf{N} . Next we choose a *polarisation type*, i.e. an $O(L)$ -orbit of a primitive vector $h \in L$. Viehweg's theory gives us the existence of a moduli space $\mathcal{M}_{n,\mathbf{N},h}$ parametrising polarised irreducible symplectic manifolds (X, \mathcal{L}) of dimension $2n$ with the chosen Beauville lattice, Fujiki invariant and polarisation type. This is a quasi-projective variety and can be constructed as a GIT quotient as in the K3 case, the only difference being that we must here invoke Matsusaka's big theorem [87] and a result of Kollár and Matsusaka [73] to be guaranteed a uniform bound N_0 such that $\mathcal{L}^{\otimes N_0}$ is very ample for all pairs (X, \mathcal{L}) .

Although there is not a Torelli theorem as in the K3 case, these moduli spaces are still related to quotients of homogeneous domains of type IV. Let Ω_L be the period domain defined by the lattice L and let $L_h = h^\perp$. This is a lattice of signature $(2, \text{rk}(L) - 3)$ and defines a homogeneous domain Ω_{L_h} of type IV. Let $O(L, h)$ be the stabiliser of h in $O(L)$. This can be considered as a subgroup of $O(L_h)$. The domain Ω_{L_h} has two connected components, of which we choose one, which we denote by \mathcal{D}_{L_h} . Again $O^+(L, h)$, the subgroup of $O(L, h)$ of real spinor norm 1, is the subgroup fixing the components of Ω_{L_h} .

Theorem 3.8. *For every component $\mathcal{M}'_{n,\mathbf{N},h}$ of the moduli space $\mathcal{M}_{n,\mathbf{N},h}$ there exists a finite to one dominant morphism*

$$\psi: \mathcal{M}'_{n,\mathbf{N},h} \rightarrow O^+(L, h) \backslash \mathcal{D}_{L_h}.$$

The proof of this is analogous to the proof of Theorem 2.19. There are, however, differences compared to the K3 case. In general, ψ will not be injective (see the discussion below). There is also a difference concerning the image of ψ . In the K3 case we know that a big and nef line bundle is ample if and only if it has positive degree on the nodal curves. Hence it is necessary and sufficient to remove the hyperplanes orthogonal to the nodal classes. So far, no complete analogue is known for the higher dimensional case, but Hassett and Tschinkel have proved partial results for $n = 2$ in [58], and more precise results in a special case in [59].

Nevertheless, Theorem 3.8 is enough to prove results on the Kodaira dimension of moduli spaces of polarised irreducible symplectic manifolds. This was done in [50], [51]: see Theorem 6.2 and Theorem 6.3 below.

Very recent work of Verbitsky [127] and Markman [86] improves Theorem 3.8, by using a polarised analogue of the monodromy group $\text{Mon}^2(X) \subset O(H^2(X), \mathbb{Z})$. Let H be an ample divisor on X . We call an element in $f \in \text{Mon}(X)$ a *polarised parallel transport operator of the pair* (X, H) if it is a parallel transport operator for a family $\pi: \mathcal{X} \rightarrow B$ with base point $b_0 \in B$ and isomorphism $\alpha: X \rightarrow \mathcal{X}_{b_0}$ which fixes $c_1(H)$, such that there exists a flat section h of $R^2\pi_*\mathbb{Z}$ with $h(b_0) = \alpha_*(c_1(H))$ and

$h(b)$ an ample class in $H^2(X_b, \mathbb{Z})$ for all $b \in B$. These operators define a subgroup $\text{Mon}^2(X, H) \subset O(H^2(X), \mathbb{Z})$. It was shown by Markman [86, Proposition 1.9] that $\text{Mon}^2(X, H)$ is the stabiliser of $c_1(H)$ in $\text{Mon}^2(X)$. Given a marking $\phi: H^2(X, \mathbb{Z}) \rightarrow L$ this defines a subgroup

$$(3.9) \quad \Gamma = \phi(\text{Mon}^2(X, H)) \subset O(L, h),$$

which can be shown to be independent of the marking ϕ : see [86, Section 7.1].

Let $\mathcal{M}'_{n, N, h}$ be a component of the moduli space of polarised irreducible symplectic manifolds with fixed numerical type and polarisation type. Given an element (X, H) in this component one thus obtains a group $\Gamma \subset O(L, h)$ as above, which is also independent of the chosen pair (X, H) by the results of [86, Section 7.1]. In fact $\Gamma \subset O^+(L, h)$, as monodromy operators are obviously orientation-preserving: see [86, Section 1.2]. Thus Γ acts on the homogeneous domain $\mathcal{D}_{L, h}$.

Theorem 3.10. *The map ψ from Theorem 3.8 lifts to an open immersion*

$$\tilde{\psi}: \mathcal{M}'_{n, N, h} \rightarrow \Gamma \backslash \mathcal{D}_{L, h},$$

where Γ is as in Equation (3.9).

Proof. It is easy to see that the map $\psi: \mathcal{M}'_{n, N, h} \rightarrow O^+(L, h) \backslash \mathcal{D}_{L, h}$ lifts to a map $\tilde{\psi}: \mathcal{M}'_{n, N, h} \rightarrow \Gamma \backslash \mathcal{D}_{L, h}$ (see the beginning of the proof of [50, Theorem 2.3]). The hard part is the injectivity of $\tilde{\psi}$ and this is where the Torelli theorem for irreducible symplectic manifolds is used. For details we refer the reader to [86]. □

Remark 3.11. We note that in general the projective group $\Gamma/\pm 1$ is a proper subgroup of $O^+(L, h)/\pm 1$ and thus Theorem 3.10 is a substantial improvement of Theorem 3.8.

In the case of irreducible symplectic manifolds of $K3^{[n]}$ type this can be made explicit. For an even lattice L we define $\text{Ref}(L)$ to be the subgroup generated by -2 -reflections and the negative of $+2$ -reflections. This is a subgroup of $O^+(L)$. If X is an irreducible symplectic manifold of $K3^{[n]}$ type, then we define $\text{Ref}(X)$ accordingly.

Theorem 3.12. (Markman) *If X is a deformation $K3^{[n]}$ manifold then*

$$\text{Mon}^2(X) = \text{Ref}(X).$$

Proof. This is proved in [83, Theorem 1.2]. □

For a lattice L we let

$$\widehat{O}(L) = \{g \in O(L) \mid g|_{L \vee L} = \pm \text{id}_{L \vee L}\}$$

and given an element $h \in L$ we set $\widehat{O}(L, h) = \{g \in \widehat{O}(L) \mid g(h) = h\}$. Recall the convention of Equation (2.15) and that by Proposition 3.2(i) the Beauville lattice is $L_{K3, 2n-2}$.

It then follows from Theorem 3.12 in conjunction with Kneser’s result [70, Satz 4] that

$$(3.13) \quad \text{Ref}(L_{K3,2n-2}) = \widehat{O}^+(L_{K3,2n-2}).$$

Combining this with Theorem 3.10 we thus obtain

Theorem 3.14. *Let \mathcal{M}'_h be an irreducible component of the moduli space of polarised deformation $K3^{[n]}$ manifolds. Then the map ψ of Theorem 3.8 factors through the finite cover $\widehat{O}^+(L_{K3,2n-2}, h) \backslash \mathcal{D}_{L_h} \rightarrow O^+(L_{K3,2n-2}, h) \backslash \mathcal{D}_{L_h}$ that is, there is a commutative diagram*

$$\begin{array}{ccc} \mathcal{M}'_h & \xrightarrow{\tilde{\psi}} & \widehat{O}^+(L_{K3,2n-2}, h) \backslash \mathcal{D}_{L_h} \\ & \searrow \psi & \downarrow \\ & & O^+(L_{K3,2n-2}, h) \backslash \mathcal{D}_{L_h}. \end{array}$$

Moreover the map $\tilde{\psi}$ is an open immersion.

Remark 3.15. In [50, Proposition 2.3] we stated that the map ψ lifts to the quotient $\widetilde{O}^+(L_{K3,2n-2}, h) \backslash \mathcal{D}_{L_h}$. This is not correct since, contrary to what was said in the proof, the projective groups $\widetilde{O}^+(L_{K3,2n-2}, h) / \pm 1$ and $\widehat{O}^+(L_{K3,2n-2}, h) / \pm 1$ are not identical if $n > 2$. In that case, in fact, $\widetilde{O}^+(L_{K3,2n-2}, h) / \pm 1$ is an index 2 subgroup of $\widehat{O}^+(L_{K3,2n-2}, h) / \pm 1$. If, however, $n = 2$, then the two groups coincide since $\widetilde{O}^+(L_{K3,2}) = \widehat{O}^+(L_{K3,2})$ and thus the results of [50] are not affected by this error.

Remark 3.16. Theorem 3.14 gives an affirmative answer to [50, Question 2.6] (with the correct group).

Remark 3.17. The results discussed so far do not give an answer to the question whether moduli spaces of polarised irreducible symplectic manifolds of given deformation type and given type of polarisation are always connected. Apostolov [1] has obtained some results on this. For example in the Hilb^[n] case these moduli spaces are always connected for $n = 2$ (both in the split and the non-split case), but in general there can be more than one component.

4. Projective models

Besides the abstract theory of moduli spaces there is a vast literature which deals with concrete geometric descriptions of K3 surfaces, and to a much lesser degree, also of irreducible symplectic manifolds of higher dimension. The easiest example is degree 4 surfaces in \mathbb{P}^3 . Any smooth quartic surface is a K3 surface and counting parameters one obtains a family of dimension $34 - 15 = 19$, because 34 is the number of quartics and 15 is the dimension of $\text{PGL}(4, \mathbb{C})$. This argument shows that the moduli space \mathcal{F}_4 of polarised K3 surfaces of degree 4 is unirational. The same approach yields unirationality for degrees $2d = 2, 6$ and 8 , as these correspond

to double covers of the projective plane branched along a sextic curve, complete intersections of type (2,3) in \mathbb{P}^4 , and complete intersections of type (2,2,2) in \mathbb{P}^5 respectively.

In general it can be very hard to decide whether a moduli space of polarised K3 surfaces of low degree is unirational or not. Mukai ([89], [90], [91], [92], [93]) has contributed most significantly to this problem. So far there are three approaches to proving unirationality.

- (1) Describing the K3 surfaces as complete intersections in homogeneous spaces (this can be used for $1 \leq d \leq 9$, $d = 11, 12, 17, 19$).
- (2) Using non-abelian Brill-Noether theory of vector bundles over algebraic curves (here one obtains results for $d = 6, 8, 10, 16$).
- (3) Using specific geometric constructions for certain degrees ($d = 11, 12, 15, 19$). An example is Mukai's most recent work ([93]) for $d = 15$ where he describes K3 surfaces of degree 30 as complete intersections in a certain rank 10 vector bundle on the Ellingsrud-Piene-Strømme moduli space of twisted cubics.

There are only a few results about rationality for these cases. Shepherd-Barron proves rationality for the cases $d = 3$ in [121] and $d = 9$ in [120].

For a discussion of low degree cases we also refer the reader to Voisin's Bourbaki exposé [130]. One can summarise the results as follows.

Theorem 4.1. *The moduli spaces \mathcal{F}_{2d} of polarised K3 surfaces of degree $2d$ are unirational for $1 \leq d \leq 12$ and $d = 15, 16, 17, 19$.*

Some other special moduli spaces related to K3 surfaces have also been studied. Perhaps the most notable of these is the moduli space of Enriques surfaces, which is a special case of the lattice polarised K3 moduli spaces given in (2.18). This approach was first seen in Horikawa's announcement [62] of a Torelli theorem for Enriques surfaces. It was shown by Kondo [75] that the moduli space of Enriques surfaces is rational.

It is also natural in this context to consider moduli of K3 surfaces with automorphisms. There is an extensive literature on such surfaces, much of it touching on moduli problems: for example [37] and the recent work of Ma and Yoshikawa [82].

Many of these moduli spaces turn out to be unirational or even rational, typically because the symmetry exhibits the K3 surfaces as covers of \mathbb{P}^2 and the family of possible ramification curves can be parametrised.

Some moduli of K3 surfaces with extra structure can be described as ball quotients: see for example [27]. Again there is an extensive literature on this subject. For a guide, we refer the reader to the survey [28] by Dolgachev and Kondo. In the introduction to [28] it is conjectured that all Deligne-Mostow arithmetic complex ball quotients are moduli spaces of K3 surfaces.

Much less is known in the case of irreducible symplectic manifolds, but some cases have been studied.

Example 4.2. A classical case is the Fano variety of lines contained in a cubic fourfold, which was studied in detail by Voisin [129]. These are varieties of $K3^{[2]}$ type. In our terminology this corresponds to the degree 6 non-split case and the lattice L_h orthogonal to the polarisation vector is isomorphic to $2U \oplus 2E_8(-1) \oplus A_2(-1)$.

Example 4.3. O'Grady studied double covers of Eisenbud-Popescu-Walter sextics in [101]. This is the case of split polarisation of minimal degree (degree 2) for the $K3^{[2]}$ -type. The lattice L_h is $2U \oplus 2E_8(-1) \oplus \langle -2 \rangle \oplus \langle -2 \rangle$.

Example 4.4. Iliev and Ranestad ([65], [66]) have shown that the variety of sums of powers $VSP(F, 10)$ of presentations of a general cubic form F in 6 variables as a sum of 10 cubes is an irreducible symplectic 4-fold. These are deformations of length 2 Hilbert schemes of K3 surfaces with a degree 14 polarisation. The precise nature of the polarisation of the irreducible symplectic manifold is unknown.

Example 4.5. Debarre and Voisin ([25]) have constructed examples in the Grassmannian $Gr(6, V)$ where V is a 10-dimensional complex vector space. Starting with a sufficiently general form $\sigma: \wedge^3 V \rightarrow \mathbb{C}$ they show that the subspace of $Gr(6, V)$ consisting of 6-planes L such that $\sigma|_{\wedge^3 L} = 0$ is an irreducible symplectic fourfold of $K3^{[2]}$ -type. This defines a 20-dimensional family with polarisation of non-split Beauville degree $2d = 22$: the lattice L_h is $2U \oplus 2E_8(-1) \oplus \begin{pmatrix} -2 & 1 \\ 1 & -6 \end{pmatrix}$.

Many authors have asked about the construction of geometrically meaningful compactifications of moduli spaces of polarised K3 surfaces. There are few general results known about this. For small degree, some results can be found in [118], [119], [123] and [115]. A partial compactification is discussed in [34] and there is an approach via log geometry in [104].

5. Compactifications

The spaces \mathcal{F}_{2d} (defined by (2.16) above) and the other quotients of period domains described in Section 4 are complex analytic spaces by construction. We observed above that they are in fact quasi-projective varieties by the results of Baily and Borel, and the GIT moduli spaces of polarised K3 surfaces and of irreducible symplectic manifolds are quasi-projective by the general results of Viehweg. Nevertheless, we require projective models and preferably smooth, or nearly smooth, models also.

In this section we describe the most commonly used compactifications and we give some results about the singularities that arise. We begin by describing the class of spaces we wish to compactify.

5.1. Modular varieties of orthogonal type

As usual we let L be an integral lattice of signature $(2, n)$, $n \geq 3$, and consider the symmetric space

$$(5.1) \quad \mathcal{D}_L = \{x \in \mathbb{P}(L \otimes \mathbb{C}) \mid (x, x) = 0, (x, \bar{x}) > 0\}^+$$

where the superscript $+$ denotes a choice of one of the two connected components of Ω_L . We let Γ be a subgroup of finite index in $O^+(L)$. Any such Γ acts properly discontinuously on \mathcal{D}_L , but in general there are elements of finite order in Γ and they have fixed points in \mathcal{D}_L .

The quotient

$$(5.2) \quad \mathcal{F}_L(\Gamma) = \Gamma \backslash \mathcal{D}_L$$

is called a *modular variety of orthogonal type* or *orthogonal modular variety*. In particular it is a locally symmetric variety, i.e. a variety that is the quotient of a symmetric space by a discrete group of automorphisms. It is not compact and is by its construction a complex analytic space: if Γ is torsion-free it is a complex manifold. In fact it is a quasi-projective variety by [5].

Some particularly important examples of orthogonal modular varieties are:

- (a) the moduli spaces of polarised K3 surfaces (the signature is $(2,19)$);
- (b) the moduli spaces of lattice-polarised K3 surfaces (signature $(2, n)$, with $n < 19$);
- (c) the moduli spaces of polarised abelian or Kummer surfaces (signature $(2,3)$);
- (d) the moduli space of Enriques surfaces (signature $(2,10)$);
- (e) quotients of the period domains of polarised irreducible symplectic varieties (signature $(2,4)$, $(2,5)$, $(2,20)$ and $(2,21)$ in the known cases).

For nearly all the orthogonal modular varieties $\mathcal{F}_L(\Gamma)$ that occur in this article, Γ is not torsion-free. The fixed points can lead to singularities of $\mathcal{F}_L(\Gamma)$. Since the stabiliser of any point of \mathcal{D}_L is finite, the singularities are finite quotient singularities: that is, locally analytically they are isomorphic to a quotient of \mathbb{C}^n by a finite subgroup G of $GL(n, \mathbb{C})$. They are not arbitrary finite quotient singularities, though, and we give some details about them in Section 5.6.

The quotient \mathbb{C}^n/G may in fact be smooth, however. This happens, by a result of Chevalley [22], if and only if G is generated by quasi-reflections (see Definition 5.19). More importantly for us, the ramification divisors of $\mathcal{D}_L \rightarrow \mathcal{F}_L(\Gamma)$ are precisely the fixed loci of elements of Γ acting as quasi-reflections on the tangent space. In fact the only quasi-reflections that occur for orthogonal modular varieties are reflections. (Let us emphasise that here we are discussing the action of an element of G , not of Γ : being a reflection on the tangent space to \mathcal{D}_L at a fixed point is not the same as being a reflection as an element of $O(L)$.)

The existence of elements acting as reflections, and thus of ramification divisors, is a significant feature of orthogonal modular varieties. For Siegel modular varieties

(the symplectic group rather than the orthogonal group) there are no ramification divisors, except in the case of Siegel modular 3-folds. Siegel modular 3-folds, however, may also be regarded as orthogonal modular varieties because of the isogeny between Sp_2 and $SO(2,3)$.

Differential forms on $\mathcal{F}_L(\Gamma)$ may be interpreted as modular forms for Γ : see Section 6.1 for more details. Therefore arithmetic information (modular forms) may be used to obtain geometric information about $\mathcal{F}_L(\Gamma)$. In particular we can use modular forms to decide whether $\mathcal{F}_L(\Gamma)$ is of general type, or more generally to try to determine its Kodaira dimension. If Y is a connected smooth projective variety of dimension n , the *Kodaira dimension* $\kappa(Y)$ of Y is defined by

$$\kappa(Y) = \text{tr.deg} \left(\bigoplus_{k \geq 0} H^0(Y, kK_Y) \right) - 1,$$

or $-\infty$ if $H^0(Y, kK_Y) = 0$ for all $k > 0$. Thus $h^0(Y, kK_Y) \sim k^{\kappa(Y)}$ for k sufficiently divisible. The possible values of $\kappa(Y)$ are $-\infty, 0, 1, \dots, n = \dim Y$, and we say Y is of *general type* if $\kappa(Y) = n$. Kodaira dimension is a bimeromorphic invariant so it makes sense to extend the definition to arbitrary irreducible quasi-projective varieties X by putting $\kappa(X) = \kappa(\tilde{X})$ for \tilde{X} a desingularisation of a compactification of X .

With this in mind, we now turn to describing some algebraic compactifications of $\mathcal{F}_L(\Gamma)$, and the singularities that can occur. Much further information about compactifications of locally symmetric varieties (not all algebraic) may be found in the book [16], especially in [16, Part III], and the references there. However, the emphasis there is on the geometry and topology of symmetric and locally symmetric spaces as real manifolds.

We are interested in two kinds of compactification: the Baily-Borel compactification $\mathcal{F}_L(\Gamma)^*$ and the toroidal compactifications $\overline{\mathcal{F}_L(\Gamma)}$. We shall describe the construction and some of the properties of each.

Remark 5.3. The constructions can also be made if Γ is an arithmetic subgroup of $O^+(L \otimes \mathbb{Q})$; that is, if $\Gamma < O(L \otimes \mathbb{Q})$ and $\Gamma \cap O^+(L)$ is of finite index in both Γ and $O^+(L)$.

In some important ways the generalisation to arithmetic subgroups does not change things much. If we are willing to change the lattice, by Proposition 5.4 we can always assume that Γ is contained in $O(L)$, so that we do not need rational entries in the matrices. In particular the results of Sections 5.5–5.7 still hold.

Proposition 5.4. *If $\Gamma < O(L \otimes \mathbb{Q})$ is arithmetic then there exists a lattice $M \subset L \otimes \mathbb{Q}$ such that $\Gamma < O(M)$.*

Proof. Let us consider $g(L)$ for all $g \in \Gamma$. The index $[\Gamma : \Gamma \cap O(L)]$ is finite, therefore the number of different copies $g(L)$ of L is finite. Therefore the \mathbb{Z} -module generated by the union of all $g(L)$ ($g \in \Gamma$) is finitely generated. Denote this lattice by M . Then Γ is a subgroup of $O(M)$ by the definition of M , and the quadratic form on M is

induced by the quadratic form on L . If the quadratic form is not even integral, then we can make it integral taking a renormalisation by an integral constant c , i.e. we set $(u, v)_M = c(u, v)_L$. Doing so does not change the orthogonal group. It follows that Γ can be considered as a subgroup of $O(M)$ for some even integral lattice M . \square

Notice, though, that the renormalisation by c does change the stable orthogonal group.

5.2. The Baily-Borel compactification

The Baily-Borel compactification, which in this context is often also referred to as the Satake compactification, can be defined very quickly as $\text{Proj} \bigoplus M_k(\Gamma, 1)$, where $M_k(\Gamma, 1)$ denotes the space of weight k modular forms with trivial character: see Definition 6.4. A priori, however, it is not clear that the ring of modular forms is finitely generated, nor that the modular forms separate points of $\mathcal{F}_L(\Gamma)$. Nor does that description immediately give a picture of the boundary $\mathcal{F}_L(\Gamma)^* \setminus \mathcal{F}_L(\Gamma)$. Instead the approach of Baily and Borel is to synthesise $\mathcal{F}_L(\Gamma)$ by topological and analytic methods, adding boundary components, and to show that the resulting space is a projective variety. Full details are given in [5], and a more detailed sketch than we give here may also be found in [16].

By writing \mathcal{D}_L in the form (5.1) we have exhibited it as a Hermitian domain of type IV. As a Riemannian domain, $\mathcal{D}_L \cong \text{SO}_0(2, n) / \text{SO}(2) \times \text{SO}(n)$, where $\text{SO}_0(2, n)$ is the identity component. The Baily-Borel compactification is defined for any Hermitian symmetric space $\mathcal{D} = G/K$ (instead of \mathcal{D}_L) and for any quotient $X = \Gamma \backslash \mathcal{D}$ of \mathcal{D} by an arithmetic group Γ . We describe the construction in general but we keep in mind the case $\mathcal{D} = \mathcal{D}_L$ as above.

An irreducible symmetric space G/K is Hermitian if and only if the centre of the maximal compact subgroup K has positive dimension: this explains why we consider only lattices of signature $(2, n)$, because $\text{SO}(m) \times \text{SO}(n)$ has discrete centre unless $n = 2$ or $m = 2$. Any Hermitian symmetric space of noncompact type can be embedded as a bounded symmetric domain in the holomorphic tangent space $T_K \mathcal{D}$ (the Harish-Chandra embedding). For these facts see [16, Prop. I.5.9] or [60].

The Baily-Borel compactification \mathcal{D}^{BB} of \mathcal{D} is simply the closure of \mathcal{D} in the Harish-Chandra embedding, which in this case is the closure $\overline{\mathcal{D}}_L$ of \mathcal{D}_L in $\mathbb{P}(L \otimes \mathbb{C})$; this in turn is contained in the compact dual, which in this case is the quadric

$$\check{\mathcal{D}} = \{x \in \mathbb{P}(L \otimes \mathbb{C}) \mid (x, x) = 0\}.$$

The tangent space to G/K at K is identified, as a complex manifold, with an open subset of $\check{\mathcal{D}}$.

A subset of \mathcal{D}^{BB} is called a *boundary component* if it is an analytic arc component: that is, an equivalence class under the relation $x \sim y$ if there exist finitely many holomorphic maps $f_i: \Delta = \{z \in \mathbb{C} \mid |z| < 1\} \rightarrow \mathcal{D}^{\text{BB}}$ such that $x \in f_1(\Delta)$, $y \in f_k(\Delta)$ and $f_i(\Delta) \cap f_{i+1}(\Delta) \neq \emptyset$ for $1 \leq i < k$. The Baily-Borel compactification \mathcal{D}^{BB}

decomposes as a disjoint union of boundary components F_p , which are themselves symmetric spaces associated with certain parabolic subgroups p of G :

$$(5.5) \quad \mathcal{D}^{\text{BB}} = \mathcal{D} \amalg \coprod_P F_p.$$

Not all parabolic subgroups occur, but only those associated with certain collections of strongly orthogonal roots. See [16, Section I.5] for precise details.

By construction, G acts on \mathcal{D}^{BB} . The normaliser of F_p ,

$$(5.6) \quad \mathcal{N}(F_p) = \{g \in G \mid g(F_p) = F_p\}$$

is a maximal parabolic subgroup of G (the parabolic subgroup P is in general not maximal). We shall later also need to consider the centraliser

$$(5.7) \quad \mathcal{Z}(F_p) = \{g \in G \mid g|_{F_p} = \text{id}\}.$$

To construct the Baily-Borel compactification of X – in our case, of $\mathcal{F}_L(\Gamma) = \Gamma \backslash \mathcal{D}_L$ – one must first restrict to rational boundary components.

Definition 5.8. A boundary component F is called a *rational boundary component* if

- (i) the normaliser $\mathcal{N}(F)$ of F in G is a parabolic subgroup defined over \mathbb{Q} , and
- (ii) the centraliser $\mathcal{Z}(F)$ contains a cocompact subgroup, normal in $\mathcal{N}(F)$, which is an algebraic subgroup defined over \mathbb{Q} .

A boundary component satisfying (i) is called a weakly rational boundary component. It is shown in [5, Theorem 3.7] that for the Baily-Borel compactification (ii) follows from (i), so that weakly rational boundary components are automatically rational.

G acts on \mathcal{D}^{BB} but of course does not preserve rational boundary components. However Γ , being an arithmetic subgroup, does take rational boundary components to rational boundary components, so it acts on $\mathcal{D}^* := \mathcal{D} \amalg \coprod_{F_p \text{ rational}} F_p$. The effect of condition (ii) is that $\Gamma_p = \Gamma \cap \mathcal{N}(F_p)$ is again a discrete group. Moreover, F_p is again a Hermitian symmetric space and Γ_p is an arithmetic group acting on F_p .

We obtain the Baily-Borel compactification $(\Gamma \backslash \mathcal{D})^*$ by taking the quotient of \mathcal{D}^* by the action of Γ . Each boundary component $\Gamma_p \backslash F_p$ has the structure of a complex analytic space and it is shown in [5] that these structures can be glued together to give an analytic structure on $(\Gamma \backslash \mathcal{D})^*$ extending the analytic structure on $\Gamma \backslash \mathcal{D}$.

It is at this stage that modular forms enter the picture. Each boundary component is an analytic space by construction but to show that their union is also an analytic space one must exhibit local analytic functions, and to show that the resulting space is projective we need analytic functions that separate points. Baily and Borel do this by using the Siegel domain realisation of \mathcal{D} over a boundary, which we describe below (Section 8.2) for the cases we need here. For a more general description, see [114]. In these coordinates one may write down suitable series (Poincaré-Eisenstein series) that define modular forms having the required properties.

Theorem 5.9. *The Baily-Borel compactification $(\Gamma \backslash \mathcal{D})^*$ is an irreducible normal projective variety over \mathbb{C} . It contains $\Gamma \backslash \mathcal{D}$ (which in our case is $\mathcal{F}_L(\Gamma)$) as a Zariski-open subset, and may be decomposed as*

$$(5.10) \quad (\Gamma \backslash \mathcal{D})^* = \Gamma \backslash \mathcal{D} \amalg \coprod_P \Gamma_P \backslash X_P,$$

where P runs through representatives of Γ -equivalence classes of parabolic subgroups determining rational boundary components.

Although we defined rational boundary components in terms of \mathcal{D}^{BB} , they determine, according to condition (i) above, rational maximal parabolic subgroups of G . If G is simple, which will always be the case for us, the boundary components of \mathcal{D}^{BB} correspond precisely to the maximal real parabolic subgroups: see [3, III.3, Proposition 3.9].

At least for the classical groups, the rational maximal parabolic subgroups can be described in combinatorial terms. In the case of $O(L)$ with L of signature $(2, n)$ they are the stabilisers of isotropic subspaces of $L \otimes \mathbb{Q}$. Because of the signature, such spaces have dimension 2 or 1 (or 0, corresponding to the “boundary” component $\Gamma \backslash \mathcal{D}_L$). Therefore we obtain the following description of $\mathcal{F}_L(\Gamma)^*$.

Theorem 5.11. *$\mathcal{F}_L(\Gamma)^*$ decomposes into boundary components as*

$$(5.12) \quad \mathcal{F}_L(\Gamma)^* = \mathcal{F}_L(\Gamma) \amalg \coprod_{\Pi} X_{\Pi} \amalg \coprod_{\ell} Q_{\ell},$$

where ℓ and Π run through representatives of the finitely many Γ -orbits of isotropic lines and isotropic planes in $L \otimes \mathbb{Q}$ respectively. Each X_{Π} is a modular curve, each Q_{ℓ} is a point, and Q_{ℓ} is contained in the closure of X_{Π} if and only if the representatives may be chosen so that $\ell \subset \Pi$.

X_{Π} and Q_{ℓ} are usually referred to as 1-dimensional and 0-dimensional boundary components, or corank 1 and corank-2 boundary components. The boundary components of the Baily-Borel compactification are also known as the *cusps*.

5.3. Toroidal compactifications

Toroidal compactifications in general are described in the book [3]. They are made by adding a divisor at each cusp. Locally in the analytic topology near a cusp, the toroidal compactification is a quotient of an open part of a toric variety over the cusp: this variety is determined by a choice of admissible fan in a suitable cone, and the choices must be made so as to be compatible with inclusions among the closures of the Baily-Borel boundary components. A summary may be found in [3, III.5].

The case we are concerned with, of $O(2, n)$, is simpler than the general case because only the 0-dimensional cusps need any attention. However, we shall begin by describing the general theory, starting with $\mathcal{D} = G/K$ and an action of an arithmetic group Γ .

Let F be a boundary component: we may as well assume immediately that it is a rational boundary component. In general one has a description of \mathcal{D} as a Siegel domain, an analytic open subset inside

$$(5.13) \quad \mathcal{D}(F) := F \times V(F) \times U(F)_{\mathbb{C}}.$$

In this decomposition, $U(F)$ is the centre of the unipotent radical $W(F)$ of $\mathcal{N}(F)$, the normaliser of F in G , and $V(F) \cong W(F)/U(F)$ is an abelian Lie group. This is not a holomorphic decomposition ($V(F)$ does not have a natural complex structure) but $U(F)_{\mathbb{C}}$ acts holomorphically on $\mathcal{D}(F)$ and in the diagram

$$\begin{array}{ccc}
 \mathcal{D}(F) & & \\
 \pi_F \downarrow & \searrow \pi'_F & \\
 & \mathcal{D}(F)' & = \mathcal{D}(F)/U(F)_{\mathbb{C}} \\
 & \swarrow p_F & \\
 F & &
 \end{array}$$

all the maps are holomorphic.

Now \mathcal{D} is given by a tube domain condition: there is a cone $C(F) \subset U(F)$ such that

$$(5.14) \quad \mathcal{D} = \{x \in \mathcal{D}(F) \mid \text{Im}(\text{pr}_U(x)) \in C(F)\}.$$

where $\text{pr}_U: \mathcal{D}(F) \rightarrow U(F)_{\mathbb{C}}$ is the projection map from the decomposition (5.13): this is a holomorphic map, even though (5.13) is not a holomorphic product decomposition.

In fact, there is a holomorphic product decomposition of $\mathcal{D}(F)$ which is (perhaps confusingly) similar:

$$(5.15) \quad \mathcal{D}(F) \cong U(F)_{\mathbb{C}} \times \mathbb{C}^k \times F,$$

where, of course, $k = \frac{1}{2} \dim_{\mathbb{R}} V(F)$ but \mathbb{C}^k is not naturally identified with $V(F)$.

Denoting the map $x \in \mathcal{D}(F) \mapsto \text{Im}(\text{pr}_U(x)) \in U(F)$ by ϕ_F , as in [3], we have the diagram

$$\begin{array}{ccccc}
 C(F) & \subset & U(F) & & \\
 \uparrow \phi_F & & \uparrow \phi_F & & \\
 \mathcal{D} & \subset & \mathcal{D}(F) & \subset & \check{\mathcal{D}} \\
 \downarrow \pi_F & \searrow & \downarrow \pi'_F & & \\
 & & \mathcal{D}(F)' & & \\
 & \swarrow p_F & & & \\
 F & & & &
 \end{array}$$

in which $\pi'_F: \mathcal{D}(F) \rightarrow \mathcal{D}(F)'$ and $p_F: \mathcal{D}(F)' \rightarrow F$ are principal homogeneous spaces for $U(F)_{\mathbb{C}}$ and $V(F)$ respectively.

When Γ acts, the group that acts on $\mathcal{D}(F)$ and on \mathcal{D} is $N(F)_{\mathbb{Z}} = \Gamma \cap N(F)$, which is a discrete group because F is a rational boundary component. So, looking at the action of $U(F)_{\mathbb{Z}} = \Gamma \cap U(F)$, we get a principal fibre bundle

$$(5.16) \quad \mathcal{D}(F)/U(F)_{\mathbb{Z}} \longrightarrow \mathcal{D}(F)'$$

whose fibre is $T(F) = U(F)_{\mathbb{C}}/U(F)_{\mathbb{Z}}$, an algebraic torus over \mathbb{C} .

Toroidal compactification proceeds by replacing this torus with a toric variety $X_{\Sigma(F)}$ and taking the closure of $\mathcal{D}/U(F)_{\mathbb{Z}}$ in the $X_{\Sigma(F)}$ -bundle over $\mathcal{D}(F)'$ that results. Doing this for each cusp F separately one can then take the quotients of each such $X_{\Sigma(F)}$ by $N(F)_{\mathbb{Z}}$ and (under suitable conditions) glue the resulting pieces together by identifying the copies of \mathcal{D}/Γ contained in each one.

In this process $\Sigma(F)$ is in general very far from unique. It is a fan in $C(F)$ (more precisely, of the convex hull of the rational points of the closure of $C(F)$), i.e. a decomposition of $C(F)$ into rational polyhedral cones (the integral structure is given by the lattice $U(F)_{\mathbb{Z}} \subset U(F)$), which is required to be $N(F)_{\mathbb{Z}}$ -equivariant and locally finite, but is not itself finite except in trivial cases: thus $X_{\Sigma(F)}$ is locally Noetherian, but not Noetherian.

In general, in order for the gluing procedure to work, the fans must satisfy a compatibility condition between fans for different cusps that arises when one cusp is in the closure of another, but in the case of $O(2, n)$ the condition is automatically satisfied. The reason is that the 1-dimensional cusps have $\dim_{\mathbb{R}} U(F) = 1$ and $C(F) = \mathbb{R}_+$, and the cone decomposition is therefore unique, and trivial. At the 0-dimensional cusps, in contrast, one has $\dim_{\mathbb{R}} U(F)$, so \mathcal{D} is actually a tube domain in $\mathcal{D}(F)$: we describe this situation explicitly in Section 8.2.

At the 0-dimensional cusps, therefore, many different choices of compactification are possible. Below we shall choose one that suits our purpose.

In the end we need to take the quotient by $N(F)_{\mathbb{Z}}$, not just $U(F)_{\mathbb{Z}}$. This has two consequences. First, this is why it is necessary to choose $\Sigma(F)$ and hence $X_{\Sigma(F)}$ in such a way that $N(F)_{\mathbb{Z}}$ still acts; secondly, even if $X_{\Sigma(F)}$ is chosen to be smooth, the action of $N(F)_{\mathbb{Z}}$ may reintroduce quotient singularities into the finished toroidal compactification. It is easy to choose $X_{\Sigma(F)}$ to be smooth, by the usual method of subdivision to resolve toric singularities.

Theorem 5.17. *A suitable choice of fans $\{\Sigma(F)\}$ for rational boundary components F determines a toroidal compactification $\overline{\mathcal{D}}/\Gamma$ of \mathcal{D}/Γ . This compactification may be chosen to be projective, and to have at worst finite quotient singularities.*

Proof. The only part not described above is the assertion that the compactification may be chosen to be projective. This is shown in [3, IV.2] with the extra assumption that Γ is neat, which is harmless because one may work with a neat normal subgroup

$\Gamma' < \Gamma$ of finite index and then use the Γ/Γ' -action. See also [31, V.5] for more details in the Siegel (symplectic group) case, in a more arithmetic framework. \square

5.4. Canonical singularities

In this part, we give sufficient conditions for the moduli space or a suitable toroidal compactification of it to have canonical singularities. We outline the proof, from [46, Section 2], that orthogonal modular varieties of dimension $n \geq 9$ satisfy these conditions.

Definition 5.18. A normal complex variety X is said to have *canonical singularities* if it is \mathbb{Q} -Gorenstein and for some (hence any) resolution of singularities $f: \tilde{X} \rightarrow X$ the discrepancy $\Delta = K_{\tilde{X}} - f^*K_X$ is an effective Weil \mathbb{Q} -divisor.

Recall that X being \mathbb{Q} -Gorenstein means that for some $r \in \mathbb{N}$, if K_X is a canonical (Weil) divisor on X then rK_X is Cartier. Therefore f^*K_X makes sense: by definition it is the \mathbb{Q} -Cartier divisor $\frac{1}{r}f^*(rK_X)$.

$\Delta = \sum \alpha_i E_i$ is supported on the irreducible exceptional divisors E_i for f , so X has canonical singularities if and only if the rational numbers α_i are all non-negative. Equivalently, X has canonical singularities if and only if on any open set $U \subset X$, any pluricanonical form (i.e. section of rK_X for some r) on the smooth part of U extends holomorphically to the whole of \tilde{U} . For more detail on canonical singularities, see [112].

A point $P \in X$ is said to be a canonical singularity if some neighbourhood X_0 of P has canonical singularities.

As we saw in Section 5.1, the singularities of $\mathcal{F}_1(\Gamma)$ are finite quotient singularities, arising at the images of points of \mathcal{D} whose stabiliser in Γ is a non-trivial finite group. Any such action can be linearised locally [21]: we therefore consider the action of a finite subgroup $G < GL(n, \mathbb{C})$ on \mathbb{C}^n and the singularities of the quotient $X = \mathbb{C}^n/G$. Any element $g \in G$ can be diagonalised since it is of finite order, and the eigenvalues of g are roots of unity.

Definition 5.19. An element g is a *quasi-reflection* if exactly one of the eigenvalues is different from 1. It is a *reflection* if that eigenvalue is -1 .

For a cyclic subgroup $\langle g \rangle \subset GL(n, \mathbb{C})$ of finite order $m > 1$, we choose a primitive m th root of unity ζ (without loss of generality, $\zeta = e^{2\pi i/m}$) and we define the Reid-Tai sum

$$(5.20) \quad \Sigma(g) = \sum \left\{ \frac{\alpha_i}{m} \right\}$$

where the eigenvalues of g are ζ^{α_i} and $\{ \}$ denotes the fractional part, $0 \leq \{q\} < 1$. For convenience we set $\Sigma(1) = 1$. The usual form of the Reid-Tai criterion is the following.

Proposition 5.21. *Suppose that G is a finite subgroup of $GL(n, \mathbb{C})$ containing no quasi-reflections. Then \mathbb{C}^n/G has canonical singularities if and only if $\Sigma(g) \geq 1$ for all $g \in G$.*

This is sufficient if one wants to classify singularities, since any quotient singularity is isomorphic to a quotient singularity where there are no quasi-reflections. In our situation, the isotropy groups sometimes do contain quasi-reflections, so we want a version of the criterion that can be applied directly in that case. First, we state a lemma that will allow us to consider the elements of G one at a time.

Lemma 5.22. *Suppose $G \subset GL(n, \mathbb{C})$ is a finite group. If $\mathbb{C}^n/\langle g \rangle$ has canonical singularities for every $g \in G$, then \mathbb{C}^n/G has canonical singularities.*

Proof. Let η be a form on $(\mathbb{C}^n/G)_{\text{reg}}$ and let $\pi: \mathbb{C}^n \rightarrow \mathbb{C}^n/G$ be the quotient map. Then $\pi^*(\eta)$ is a G -invariant regular form on $\mathbb{C}^n \setminus \pi^{-1}(\mathbb{C}^n/G)_{\text{sing}}$. Since $\pi^{-1}(\mathbb{C}^n/G)_{\text{sing}}$ has codimension at least 2, the form $\pi^*(\eta)$ extends by Hartog’s theorem to a G -invariant regular form on all of \mathbb{C}^n . Now the claim follows from [124, Proposition 3.1], which says that a G -invariant form on \mathbb{C}^n extends to a desingularisation of \mathbb{C}^n/G if and only if it extends to a desingularisation of $\mathbb{C}^n/\langle g \rangle$ for every $g \in G$. \square

The converse of Lemma 5.22 is false.

Suppose that $g \in GL(n, \mathbb{C})$ is of order $m = sk$, where k is the smallest positive integer such that g^k is either a quasi-reflection or the identity. Order the eigenvectors so that the first $n - 1$ eigenvalues of g^k are equal to 1, so that the last eigenvalue is a primitive s th root of unity. We define a modified Reid-Tai sum

$$(5.23) \quad \Sigma'(g) = \left\{ \frac{sa_n}{m} \right\} + \sum \left\{ \frac{a_i}{m} \right\}$$

where again the eigenvalues of g are ζ^{a_i} , and put $\Sigma'(1) = 1$. The idea (originating in an observation of Katharina Ludwig) is that this enables us to handle the quasi-reflections correctly.

Proposition 5.24. *Suppose that G is a finite subgroup of $GL(n, \mathbb{C})$. Then \mathbb{C}^n/G has canonical singularities if $\Sigma'(g) \geq 1$ for all $g \in G$.*

Proof. If no power of g is a quasi-reflection then $s = 1$ and the usual Reid-Tai criterion (Proposition 5.21) shows that $\mathbb{C}^n/\langle g \rangle$ has canonical singularities.

Otherwise, consider g with $g^k = h$ a quasi-reflection as above. The eigenvalues of g are $\zeta^{a_1}, \dots, \zeta^{a_n}$, where ζ is a primitive m th root of unity, $\text{hcf}(s, a_n) = 1$ and $s|a_i$ for $i < n$. The group $\langle h \rangle$ is generated by quasi-reflections so $\mathbb{C}^n/\langle h \rangle \cong \mathbb{C}^n$, and we need to look at the action of the group $\langle g \rangle/\langle h \rangle$ on $\mathbb{C}^n/\langle h \rangle$. The eigenvalues of the differential of $g^l\langle h \rangle$ on $\mathbb{C}^n/\langle h \rangle$ are $\zeta^{la_1}, \dots, \zeta^{la_{n-1}}, \zeta^{sla_n}$, so

$$(5.25) \quad \Sigma(g^l\langle h \rangle) = \Sigma'(g^l) \geq 1.$$

Thus $(\mathbb{C}^n/\langle h \rangle)/\langle g\langle h \rangle \rangle \cong \mathbb{C}^n/\langle g \rangle$ has canonical singularities and the result follows by Lemma 5.22. \square

5.5. Singularities of modular varieties

We are interested primarily in the singularities of \mathcal{F}_{2d} and the other spaces mentioned in Section 4, but we may more generally consider the singularities of compactified locally symmetric varieties associated with the orthogonal group of a lattice of signature $(2, n)$. Unless n is small, it turns out that the compactification may be chosen to have canonical singularities.

Theorem 5.26. *Let L be a lattice of signature $(2, n)$ with $n \geq 9$, and let $\Gamma < O^+(L)$ be a subgroup of finite index. Then there exists a projective toroidal compactification $\overline{\mathcal{F}}_L(\Gamma)$ of $\mathcal{F}_L(\Gamma) = \Gamma \backslash \mathcal{D}_L$ such that $\overline{\mathcal{F}}_L(\Gamma)$ has canonical singularities and there are no branch divisors in the boundary. The branch divisors in $\mathcal{F}_L(\Gamma)$ arise from the fixed divisors of \pm reflections.*

There are three parts to the proof of this theorem. One must first consider the singularities of the open part, which means working out some details of the action of Γ on \mathcal{D}_L . Then there are the possible singularities over the 0-dimensional cusps: these lead to toric questions, and here one must choose the toroidal compactification appropriately. Finally, in order to deal with the 1-dimensional cusps we need a full description of the geometry there.

5.6. Singularities in the interior

Here we are interested in the singularities that arise at fixed points of the action of Γ on \mathcal{D}_L . Let $w \in L_{\mathbb{C}}$ and let $G \subset \Gamma$ be the stabiliser of $[w] \in \mathcal{D}_L$. For $[w] \in \mathcal{D}_L$ we define $\mathbb{W} = \mathbb{C}w$. Then G acts on \mathbb{W} , so $g(w) = \alpha(g)w$ for some character $\alpha: G \rightarrow \mathbb{C}^*$, and we put $G_0 = \ker \alpha$. We also put $S = (\mathbb{W} \oplus \overline{\mathbb{W}})^{\perp} \cap L$ (possibly $S = \{0\}$) and $T = S^{\perp} \subset L$. In the case of polarised K3 surfaces, S is the primitive part of the Picard lattice and T is the transcendental lattice of the surface corresponding to the period point w .

It is easy to check that $S_{\mathbb{C}} \cap T_{\mathbb{C}} = \{0\}$ and that G acts on S and on T : moreover G_0 acts trivially on $T_{\mathbb{Q}}$.

Since $G/G_0 \subset \text{Aut } \mathbb{W} \cong \mathbb{C}^*$ it is a cyclic group: we denote its order by r_w . So by the above, $\mu_{r_w} \cong G/G_0$ acts on $T_{\mathbb{Q}}$. (By μ_r we mean the group of r th roots of unity in \mathbb{C} .)

For any $r \in \mathbb{N}$ there is a unique faithful irreducible representation of μ_r over \mathbb{Q} , which we call \mathcal{V}_r . The dimension of \mathcal{V}_r is $\varphi(r)$, where φ is the Euler φ function and, by convention, $\varphi(1) = \varphi(2) = 1$. The eigenvalues of a generator of μ_r in this representation are precisely the primitive r th roots of unity: \mathcal{V}_1 is the 1-dimensional trivial representation. Note that $-\mathcal{V}_d = \mathcal{V}_d$ if d is even and $-\mathcal{V}_d = \mathcal{V}_{2d}$ if d is odd.

Using the fact that $S_{\mathbb{C}} \cap T_{\mathbb{C}} = 0$, we may check that $T_{\mathbb{Q}}$ splits as a direct sum of irreducible representations \mathcal{V}_{r_w} (in particular, $\varphi(r_w) \mid \dim T_{\mathbb{Q}}$) and that if $g \in G$ and $\alpha(g)$ is of order r (so $r \mid r_w$), then $T_{\mathbb{Q}}$ splits as a g -module into a direct sum of irreducible representations \mathcal{V}_r of dimension $\varphi(r)$.

We are interested in the action of G on the tangent space to \mathcal{D}_L . We have a natural isomorphism

$$T_{[\mathbf{w}]} \mathcal{D}_L \cong \text{Hom}(\mathbb{W}, \mathbb{W}^\perp / \mathbb{W}) =: V.$$

Suppose $g \in G$ is of order m and $\alpha(g)$ is of order r : as usual we take $\zeta = e^{2\pi i/m}$, and henceforth we think of g as an element of $GL(V)$, with eigenvalues $\zeta^{\alpha_1}, \dots, \zeta^{\alpha_n}$.

If $\varphi(r)$ is not very small, the copy of \mathcal{V}_r containing \mathbf{w} already contributes at least 1 to $\Sigma(g)$. The cases $r = 1$ and $r = 2$ are also simple.

Proposition 5.27. *Assume that $g \in G$ does not act as a quasi-reflection on V and that $\varphi(r) > 4$. Then $\Sigma(g) \geq 1$.*

Proof. As ξ runs through the m th roots of unity, $\xi^{m/r}$ runs through the r th roots of unity. We denote by $k_1, \dots, k_{\varphi(r)}$ the integers such that $0 < k_i < r$ and $(k_i, r) = 1$, in no preferred order. Without loss of generality, we assume $\alpha(g) = \zeta^{mk_2/r}$ and $\overline{\alpha(g)} = \alpha(g)^{-1} = \zeta^{mk_1/r}$, with $k_1 \equiv -k_2 \pmod r$.

One of the \mathbb{Q} -irreducible subrepresentations of g on $L_{\mathbb{C}}$ contains the eigenvector \mathbf{w} : we call this $\mathbb{V}_r^{\mathbf{w}}$ (it is the smallest g -invariant complex subspace of $L_{\mathbb{C}}$ that is defined over \mathbb{Q} and contains \mathbf{w}). It is a copy of $\mathcal{V}_r \otimes \mathbb{C}$: to distinguish it from other irreducible subrepresentations of the same type we write $\mathbb{V}_r^{\mathbf{w}} = \mathcal{V}_r^{\mathbf{w}} \otimes \mathbb{C}$.

If \mathbf{v} is an eigenvector for g with eigenvalue $\zeta^{mk_i/r}$, $i \neq 1$ (in particular $\mathbf{v} \notin \overline{\mathbb{W}}$), then $\mathbf{v} \in \mathbb{W}^\perp$ since $(\mathbf{v}, \mathbf{w}) = (g(\mathbf{v}), g(\mathbf{w})) = \zeta^{mk_i/r} \alpha(g)(\mathbf{v}, \mathbf{w})$. Therefore the eigenvalues of g on $\mathbb{V}_r^{\mathbf{w}} \cap \mathbb{W}^\perp / \mathbb{W}$ include $\zeta^{mk_i/r}$ for $i \geq 3$, so the eigenvalues on $\text{Hom}(\mathbb{W}, \mathbb{V}_r^{\mathbf{w}} \cap \mathbb{W}^\perp / \mathbb{W}) \subset V$ include $\zeta^{mk_1/r} \zeta^{mk_i/r}$ for $i \geq 3$. So

$$(5.28) \quad \Sigma(g) \geq \sum_{i=3}^{\varphi(r)} \left\{ \frac{k_1}{r} + \frac{k_i}{r} \right\} \geq 1,$$

where the last inequality is an elementary verification. □

The cases $r = 1$ and $r = 2$ are simple to deal with because one always finds two conjugate eigenvalues, which between them contribute 1 to $\Sigma(g)$.

So far we have needed no hypothesis on the dimension, but the remaining cases ($r = 3, 4, 5, 6, 8, 10$ or 12) do require such a condition because the contributions to $\Sigma(g)$ from each \mathcal{V}_r are small and we need to have enough of them. We refer to [46] for details. In the end we find

Theorem 5.29. *Assume that $g \in G$ does not act as a quasi-reflection on V and that $n \geq 6$. Then $\Sigma(g) \geq 1$.*

One must then carry out a similar analysis for quasi-reflections. One more dimension is needed to guarantee $\Sigma'(g) \geq 1$, because the a_n term does not help us. The analysis also has the corollary that the quasi-reflections in the tangent space V that arise are in this case always reflections, and moreover that the elements of Γ that they come from are themselves (up to sign) reflections, considered as elements of $O(L)$.

Corollary 5.30. *If $n \geq 7$ then $\mathcal{F}_L(\Gamma)$ has canonical singularities.*

5.7. Singularities at the cusps

We now consider the boundary $\overline{\mathcal{F}}_L(\Gamma) \setminus \mathcal{F}_L(\Gamma)$. Cusps, or boundary components in the Baily-Borel compactification, correspond to orbits of totally isotropic subspaces $E \subset L_{\mathbb{Q}}$. Since L has signature $(2, n)$, the dimension of E is 1 or 2, corresponding to dimension 0 and dimension 1 boundary components respectively.

A toroidal compactification over a cusp F coming from an isotropic subspace E corresponds to an admissible fan Σ in some cone $C(F) \subset U(F)$. We have, as in [3]

$$\mathcal{D}_L(F) := U(F)_{\mathbb{C}} \mathcal{D}_L \subset \check{\mathcal{D}}_L$$

where $\check{\mathcal{D}}_L$ is the compact dual of \mathcal{D}_L (see [3, III.2]).

The case $\dim E = 1$, that is, isotropic vectors in L , is the case of 0-dimensional cusps in the Baily-Borel compactification and leads to a purely toric problem. In this case we have

$$\mathcal{D}_L(F) \cong F \times U(F)_{\mathbb{C}} = U(F)_{\mathbb{C}}.$$

Put $M(F) = U(F)_{\mathbb{Z}}$ and define the torus $\mathbf{T}(F) = U(F)_{\mathbb{C}}/M(F)$. In general $(\mathcal{D}_L/M(F))_{\Sigma}$ is by definition the interior of the closure of $\mathcal{D}_L/M(F)$ in $\mathcal{D}_L(F)/M(F) \times_{\mathbf{T}(F)} X_{\Sigma}(F)$, i.e. in $X_{\Sigma}(F)$ in this case, where $X_{\Sigma}(F)$ is the torus embedding corresponding to the torus $\mathbf{T}(F)$ and the fan Σ . We may choose Σ so that $X_{\Sigma}(F)$ is smooth and $G(F) := N(F)_{\mathbb{Z}}/U(F)_{\mathbb{Z}}$ acts on $(\mathcal{D}_L/M(F))_{\Sigma}$: this is also implicit in [3] and explained in [31, p.173]. The toroidal compactification is locally isomorphic to $X_{\Sigma}(F)/G(F)$. Thus the problem of determining the singularities is reduced to a question about toric varieties, which is answered by Theorem 5.31, below.

We take a lattice M of dimension n and denote its dual lattice by N . A fan Σ in $N \otimes \mathbb{R}$ determines a toric variety X_{Σ} with torus $\mathbf{T} = \text{Hom}(M, \mathbb{C}^*) = N \otimes \mathbb{C}^*$.

Theorem 5.31. *Let X_{Σ} be a smooth toric variety and suppose that a finite group $G < \text{Aut}(\mathbf{T}) = \text{GL}(M)$ of torus automorphisms acts on X_{Σ} . Then X_{Σ}/G has canonical singularities.*

Proof. This is [46, Theorem 2.17]. The proof also shows (with a little modification) that there are no branch divisors contained in the boundary over 0-dimensional cusps either. □

It remains to consider the dimension 1 cusps. We consider a rank 2 totally isotropic subspace $E_{\mathbb{Q}} \subset L_{\mathbb{Q}}$, corresponding to a dimension 1 boundary component F of \mathcal{D}_L . The idea is to choose standard bases for $L_{\mathbb{Q}}$ so as to be able to identify $U(F)$, $U(F)_{\mathbb{Z}}$ and $N(F)_{\mathbb{Z}}$ explicitly, as is done in [115] for maximal K3 lattices, where $n = 19$. Then, following Kondo [74] one can analyse the group action in coordinates, using the Siegel domain realisation of \mathcal{D} associated with the given cusp. Both in [74] and in [115] there are special features that allow one to work over \mathbb{Z} , but in general one must work over \mathbb{Q} . For details we refer to [46].

6. Modular forms and Kodaira dimension

One of the main tools in the study of the geometry of the orthogonal modular varieties $\mathcal{F}_L(\Gamma)$ is the theory of modular forms with respect to an orthogonal group of type $O(2, n)$. One application is to prove that $\mathcal{F}_L(\Gamma)$ is often of general type. The methods described here were used in [46] to prove the following result.

Theorem 6.1. *The moduli space \mathcal{F}_{2d} of K3 surfaces with a polarisation of degree $2d$ is of general type for any $d > 61$ and for $d = 46, 50, 52, 54, 57, 58$ and 60 .*

If $d \geq 40$ and $d \neq 41, 44, 45$ or 47 then the Kodaira dimension of \mathcal{F}_{2d} is non-negative.

Similar methods apply to irreducible symplectic manifolds and their polarisations, discussed in Section 3.3. For deformations of length 2 Hilbert schemes of K3 surfaces with polarisation of split type (see Equation (3.6)) there is the following result, from [50].

Theorem 6.2. *The variety $\mathcal{M}_{2d}^{[2],split}$ is of general type if $d \geq 12$. Moreover its Kodaira dimension is non-negative if $d = 9$ and $d = 11$.*

For the ten-dimensional O’Grady case [99], there are again split and non-split polarisations, and a fairly complete general type result in the split case was proved in [51].

Theorem 6.3. *Let d be a positive integer not equal to 2^n with $n \geq 0$. Then every component of the moduli space of ten-dimensional polarised O’Grady varieties with split polarisation h of Beauville degree $h^2 = 2d \neq 2^{n+1}$ is of general type.*

We do not attempt to prove Theorem 6.3 here but the theory we develop in the rest of this article will give proofs (though not with full details) of Theorem 6.1 and Theorem 6.2.

6.1. Modular forms of orthogonal type

In Definition 6.4 below we follow [11]. An “affine” definition similar to the one usually given for of $SL(2)$ can be found in [40]. The *affine cone* over \mathcal{D}_L is $\mathcal{D}_L^\bullet = \{y \in L \otimes \mathbb{C} \mid x = \mathbb{C}^*y \in \mathcal{D}_L\}$.

Definition 6.4. Suppose that L has signature $(2, n)$, with $n \geq 3$. Let $k \in \mathbb{Z}$ and let $\chi: \Gamma \rightarrow \mathbb{C}^*$ be a character of a subgroup $\Gamma < O^+(L)$ of finite index. A holomorphic function $F: \mathcal{D}_L^\bullet \rightarrow \mathbb{C}$ is called a *modular form* of *weight* k and character χ for the group Γ if

$$F(tZ) = t^{-k}F(Z) \quad \forall t \in \mathbb{C}^*,$$

$$F(gZ) = \chi(g)F(Z) \quad \forall g \in \Gamma.$$

A modular form is called a *cuspidal form* if it vanishes at every cusp.

The weight as defined here is what is sometimes called *arithmetic weight*. Some authors prefer to use the geometric weight, which is k/n , normally only in contexts where $n|k$. We shall always use the arithmetic weight. One may choose a complex volume form dZ on \mathcal{D}_L such that if F is a modular form of weight mn and character \det^m for Γ then $F(dZ)^m$ is a Γ -invariant section of $mK_{\mathcal{D}_L}$: see [7] for a precise account.

If $n < 3$ then one has to add to Definition 6.4 the condition that F is holomorphic at the boundary. According to Koecher’s principle (see [4], [32], [107]) this condition is automatically fulfilled if the dimension of a maximal isotropic subspace of $L \otimes \mathbb{Q}$ is smaller than n . In particular, this is always true if $n \geq 6$.

We denote the linear spaces of modular and cusp forms of weight k and character χ by $M_k(\Gamma, \chi)$ and $S_k(\Gamma, \chi)$ respectively. If $M_k(\Gamma, \chi)$ is nonzero then one knows that $k \geq (n - 2)/2$ (see [40]). The minimal weight $k = (n - 2)/2$ is called *singular*. Modular forms of singular weight are very special. The first example of such forms for orthogonal groups was constructed in [38]. Cusp forms are possible only if $k > (n - 2)/2$. The weight $k = \dim(\mathcal{F}_L(\Gamma))$ is called *canonical* because by a lemma of Freitag

$$S_n(\Gamma, \det) \cong H^0(\tilde{\mathcal{F}}_L(\Gamma), K_{\tilde{\mathcal{F}}_L(\Gamma)}),$$

where $\tilde{\mathcal{F}}_L(\Gamma)$ is a smooth compact model of the modular variety $\mathcal{F}_L(\Gamma)$ and $K_{\tilde{\mathcal{F}}_L(\Gamma)}$ is the sheaf of canonical differential forms (see [32, Hilfssatz 2.1, Kap. 3]). Therefore we have the following important formula for the geometric genus of the modular variety:

$$(6.5) \quad p_g(\tilde{\mathcal{F}}_L(\Gamma)) = \dim S_n(\Gamma, \det).$$

Remark 6.6. Below (see Section 8.3) we describe the property of being holomorphic at the boundary (needed only if $n \leq 2$) in terms of the Fourier expansions.

Remark 6.7. In this article we usually assume that $n \geq 3$. In this case the order of any character χ in Definition 6.4 is finite according to Kazhdan’s property (T) (see [69]).

Remark 6.8. If the lattice L contains two orthogonal copies of the hyperbolic plane $U \cong \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and if its reduction modulo 2 (respectively 3) is of rank at least 6 (respectively 5) then $\tilde{O}^+(L)$ has only one non-trivial character, namely \det (see [49]). In particular the modular group $\tilde{O}^+(L_{2d})$ related to the polarised K3 surfaces has only one non-trivial character.

Remark 6.9. If $L_{2t}^{(5)} = 2U \oplus \langle -2t \rangle$, of signature $(2,3)$, then the modular forms with respect to $\tilde{SO}^+(L_{2t}^{(5)})$ coincide with Siegel modular forms with respect to the paramodular group Γ_t (see [39], [43], [53], [55]). In particular, if $t = 1$ we obtain the Siegel modular forms with respect to $Sp_2(\mathbb{Z})$. In contrast to Remark 6.8 the group $\tilde{SO}^+(L_{2t}^{(5)})$ has non-trivial characters. They were described in [44, Section 2]. One can

construct important cusp forms of the minimal possible weight 1 with non-trivial character for the full modular group $\widetilde{SO}^+(L_{2t}^{(5)}) \cong \Gamma_t/\{\pm 1\}$ for some t (see [55]).

6.2. Rational quadratic divisors

For any $v \in L \otimes \mathbb{Q}$ such that $v^2 = (v, v) < 0$ we define the *rational quadratic divisor*

$$(6.10) \quad \mathcal{D}_v = \mathcal{D}_v(L) = \{[Z] \in \mathcal{D}_L \mid (Z, v) = 0\} \cong \mathcal{D}_{v^\perp}$$

where v^\perp is an even integral lattice of signature $(2, n - 1)$. Therefore \mathcal{D}_v is also a homogeneous domain of type IV. We note that $\mathcal{D}_v(L) = \mathcal{D}_{tv}(L)$ for any $t \neq 0$. The theory of automorphic Borchers products (see [13]) gives a method of constructing automorphic forms with rational quadratic divisors. Special divisors of this type (the reflective divisors defined below) play an important role in the theory of moduli spaces.

The reflection with respect to the hyperplane defined by a non-isotropic vector r is given by

$$(6.11) \quad \sigma_r: l \mapsto l - \frac{2(l, r)}{(r, r)} r.$$

If r is primitive in L and the reflection σ_r fixes L , i.e. $\sigma_r \in O(L)$, then we say that r is a *reflective vector*, also known as a *root*. If $(r, r) = d$ we say that r is a d -vector or (if it is a root) a d -root. A 2-vector or a -2 -vector is always a root.

If $v \in L^\vee$ and $(v, v) < 0$, the divisor $\mathcal{D}_v(L)$ is called a *reflective divisor* if $\sigma_v \in O(L)$. It was proved in [46, Corollary 2.13] that for $n \geq 3$ the branch divisor of the modular projection

$$\pi_\Gamma: \mathcal{D}_L \rightarrow \Gamma \backslash \mathcal{D}_L$$

is the union of the reflective divisors with respect to Γ :

$$(6.12) \quad \text{Bdiv}(\pi_\Gamma) = \bigcup_{\mathbb{Z}r \subset L, \sigma_r \in \Gamma \cup -\Gamma} \mathcal{D}_r(L).$$

Note that here we have to allow r such that $-\sigma_r \in \Gamma$ as well as those with $\sigma_r \in \Gamma$: compare Remark 6.16 below, concerning modular forms.

6.3. Low weight cusp form trick

The next theorem, proved in [46, Theorem 1.1], is called the *low weight cusp form trick*. It plays a crucial role in the application of modular forms to moduli problems. If F is a modular form (of any weight or character) then the *divisor* $\text{div } F$ in \mathcal{D}_L is given by the equation $F(Z) = 0$: this is well-defined in view of Definition 6.4.

Theorem 6.13. *Let L be an integral lattice of signature $(2, n)$, $n \geq 9$. The modular variety $\mathcal{F}_L(\Gamma)$ is of general type if there exists a non-zero cusp form $F_\alpha \in S_\alpha(\Gamma, \chi)$ of small weight $\alpha < n$ vanishing with order at least 1 at infinity such that $\text{div } F_\alpha \geq \text{Bdiv}(\pi_\Gamma)$.*

Proof. We let $\overline{\mathcal{F}}_L(\Gamma)$ be a projective toroidal compactification of $\mathcal{F}_L(\Gamma)$ with canonical singularities and no ramification divisors at infinity, which exists by Theorem 5.26. We take a smooth projective model Y of $\mathcal{F}_L(\Gamma)$ by taking a resolution of singularities of $\overline{\mathcal{F}}_L(\Gamma)$. We want to show the existence of many pluricanonical forms on Y .

Suppose that $F_{nk} \in M_{nk}(\Gamma, \det^k)$. By choosing a 0-dimensional cusp we may realise \mathcal{D}_L as a tube domain (see Section 8.2 for details) and use this to select a holomorphic volume element dZ . Then the differential form $\Omega(F_{nk}) = F_{nk} (dZ)^k$ is Γ -invariant and therefore determines a section of the pluricanonical bundle $kK = kK_Y$ away from the branch locus of $\pi: \mathcal{D}_L \rightarrow \mathcal{F}_L(\Gamma)$ and the cusps: see [3, p. 190] (but note that weight 1 in the sense of [3] corresponds to weight n in our definition).

In general $\Omega(F_{nk})$ will not extend to a global section of kK . We distinguish three kinds of obstruction to its doing so. There are *elliptic obstructions*, arising because of singularities given by elliptic fixed points of the action of Γ ; *reflective obstructions*, arising from the ramification divisors in \mathcal{D}_L ; and *cuspidal obstructions*, arising from divisors at infinity.

In order to deal with these obstructions we consider a neat normal subgroup Γ' of Γ of finite index and set $G := \Gamma/\Gamma'$. Let $X := \mathcal{F}_L(\Gamma')$ and let $\overline{X} := \overline{\mathcal{F}}_L(\Gamma')$ be the toroidal compactification of $\mathcal{F}_L(\Gamma')$ given by the same choice of fan as for $\overline{\mathcal{F}}_L(\Gamma)$. Then \overline{X} is a smooth projective manifold with $\overline{\mathcal{F}}_L(\Gamma) = \overline{X}/G$. Let $D := \overline{X} \setminus X$ be the boundary divisor of \overline{X} . For any element $g \in G$ we define its fixed locus $\overline{X}^g := \{x \in \overline{X} \mid g(x) = x\}$ and denote its divisorial part by $\overline{X}_{(1)}^g$. Then $R := \bigcup_{g \neq 1} \overline{X}_{(1)}^g$ is the ramification divisor of the map $\pi: \overline{X} \rightarrow \overline{X}/G$.

The results of Section 5.5 (see Theorem 5.26 and Theorem 5.31) can be summarised as follows:

- (i) R does not contain a component of D ;
- (ii) the ramification index of $\pi: \overline{X} \rightarrow \overline{X}/G$ along R is 2;
- (iii) \overline{X}/G has canonical singularities.

We will now apply the low-weight cusp form trick, used for example in [32] (for Siegel modular forms), [40], [42] and [57]. The main point is to use special cusp forms. For this let the order of χ be N and assume that k is a multiple of $2N$. Then we consider forms $F_{nk}^0 \in S_{nk}(\Gamma, \det^k) = S_{nk}(\Gamma, 1)$ of the form

$$F_{nk}^0 = F_a^k F_{(n-a)k}$$

where $F_{(n-a)k} \in M_{(n-a)k}(\Gamma, 1)$ is a modular form of weight $(n-a)k \geq k$. We claim that the corresponding forms $\Omega(F_{nk}^0)$ give rise to pluricanonical forms on Y . To see this, we deal with the three kinds of obstruction in turn.

Cuspidal obstructions. By definition, $\Omega(F_{nk}^0)$ is a G -invariant holomorphic section of kK_X . Since F_a is a cusp form of weight $a < n$, the form F_{nk}^0 has zeroes of order k along the boundary D and hence extends to a G -invariant holomorphic section of $kK_{\overline{X}}$ by [3, IV.1, Theorem 1.1].

Reflective obstructions. Since $R \subset \text{div}(F_\alpha)$ by assumption, $\Omega(F_{nk}^0)$ has zeroes of order k on $R \setminus D$. By (i) above, $\Omega(F_{nk}^0)$ actually has zeroes of order k along all of R . By (ii) the form $\Omega(F_{nk}^0)$ descends to a holomorphic section of $kK_{(\bar{X}/G)_{\text{reg}}}$ where $(\bar{X}/G)_{\text{reg}}$ is the regular part of \bar{X}/G .

Elliptic obstructions. By (iii) the form $\Omega(F_{nk}^0)$ extends to a holomorphic section of kK_Y .

Therefore $F_\alpha^k M_{(n-a)k}(\Gamma, 1)$ is a subspace of $H^0(Y, kK_Y)$. The theorem now follows because by Hirzebruch-Mumford proportionality (see [94]), $\dim M_{(n-a)k}(\Gamma, 1)$ grows like k^n . □

Remark 6.14. There is another way to deal with the reflective obstructions, which works even if a cusp form with the right properties cannot be found. Among forms of very high weight there must be some that vanish along the reflective divisors, because $\dim M_k(\Gamma, 1)$ grows faster with k than the space of obstructions, which are sections in some bundles on the reflective divisors. In [48] we estimate these dimensions using Hirzebruch-Mumford proportionality. This method can be used to produce general type results even in cases where special forms constructed by quasi pull-back are not available, but if the quasi pull-back method is available it normally produces much stronger results.

6.4. Reflective modular forms

For Theorem 6.13 we used cusp forms of low weight ($k < n$) with large divisor ($\text{div } F \geq \text{Bdiv}(\pi_\Gamma)$). We construct such modular forms for the moduli spaces of polarised K3 surfaces and other holomorphic symplectic varieties in Section 8. Modular forms of high weight ($k \geq n$) with small divisor ($\text{div } F \leq \text{Bdiv}(\pi_\Gamma)$) also have applications to the theory of moduli spaces such as Theorem 6.19 below.

Definition 6.15. A modular form $F \in M_k(\Gamma, \chi)$ is called *reflective* if

$$\text{Supp}(\text{div } F) \subset \bigcup_{\mathbb{Z}\tau \subset L, \sigma_\tau \in \Gamma \cup -\Gamma} \mathcal{D}_\tau(L) = \text{Bdiv}(\pi_\Gamma).$$

We call F *strongly reflective* if the multiplicity of any irreducible component of $\text{div } F$ is equal to one.

Remark 6.16. In the definition of reflective modular forms given in [54] only the condition $\sigma_\tau \in \Gamma$ was considered. The present definition, allowing $-\sigma_\tau \in \Gamma$, is explained by equation (6.12).

Example 6.17. The most famous example of a strongly reflective modular form is the *Borchers modular form* $\Phi_{12} \in M_{12}(\mathcal{O}^+(\text{II}_{2,26}), \det)$ (see [11]). This is the unique modular form of singular weight 12 with character \det with respect to the orthogonal group $\mathcal{O}^+(\text{II}_{2,26})$ of the even unimodular lattice $\text{II}_{2,26} \cong 2U \oplus 3E_8(-1)$ of signature $(2, 26)$. The form Φ_{12} is the Kac-Weyl-Borchers denominator function of the Fake

Monster Lie algebra. For any (-2) -vector $r \in \text{II}_{2,26}$ we have

$$\Phi_{12}(\sigma_r(Z)) = \det(\sigma_r)\Phi_{12}(Z) = -\Phi_{12}(Z).$$

Therefore Φ_{12} vanishes along $\mathcal{D}_r(\text{II}_{2,26})$. According to [11] the order of vanishing is 1 and the full divisor of this modular form is the union of the mirrors of such reflections:

$$\text{div}_{\mathcal{D}(\text{II}_{2,26})} \Phi_{12} = \sum_{\substack{\pm r \in \text{II}_{2,26} \\ r^2 = -2}} \mathcal{D}_r(\text{II}_{2,26}).$$

According to Eichler’s criterion (see Lemma 7.8) all (-2) -vectors of $\text{II}_{2,26}$ constitute one $\tilde{\mathcal{O}}^+(\text{II}_{2,26})$ -orbit. In other words, the ramification divisor of the 26-dimensional modular variety $\mathcal{F}_{\text{II}_{2,26}}(\tilde{\mathcal{O}}^+(\text{II}_{2,26}))$ is irreducible.

Remark 6.18. Modular forms of weight n (known as canonical weight) have special properties. Suppose L has signature $(2, n)$ and $F \in M_n(\Gamma, \det)$. If $\sigma_r \in \Gamma$, then $F(\sigma_r(Z)) = -F(Z)$. Hence F vanishes along $\mathcal{D}_r(L)$. If $-\sigma_r \in \Gamma$, then

$$(-1)^n F(\sigma_r(Z)) = F(-\sigma_r(Z)) = \det(-\sigma_r)F(Z) = (-1)^{n+1}F(Z)$$

and F also vanishes along $\mathcal{D}_r(L)$. Therefore any Γ -modular form of canonical weight with character \det vanishes along $\text{Bdiv}(\pi_\Gamma)$.

If $S_n(\Gamma, \det) \neq 0$ then the Kodaira dimension of $\overline{\mathcal{F}}_L(\Gamma)$ is non-negative (with no restriction needed on the dimension n), because if $F_n \in S_n(\Gamma, \det)$ then by Freitag’s lemma $F_n(Z)dZ$ defines an element of $H^0(\overline{\mathcal{F}}_L(\Gamma), K_{\overline{\mathcal{F}}_L(\Gamma)})$. Therefore $p_g(\overline{\mathcal{F}}_L(\Gamma)) \geq 1$ and the plurigenera do not all vanish.

The next theorem was proved in [41] and contrasts with Theorem 6.13.

Theorem 6.19. *Suppose that L has signature $(2, n)$, with $n \geq 3$. Let $F_k \in M_k(\Gamma, \chi)$ be a strongly reflective modular form of weight k and character χ for a subgroup $\Gamma < \mathcal{O}^+(L)$ of finite index. Then*

$$(6.20) \quad \kappa(\mathcal{F}_L(\Gamma)) = -\infty$$

if $k > n$, or if k and F_k is not a cusp form. If k and F_n is a cusp form whose order of zero at infinity is at least 1 then

$$(6.21) \quad \kappa(\Gamma_\chi \backslash \mathcal{D}_L) = 0,$$

where $\Gamma_\chi = \ker(\chi \cdot \det)$ is a subgroup of Γ .

Proof. The requirement that the cusp form should have order of vanishing at least 1 is almost always satisfied: see [49].

To prove (6.20) we have to show that there are no pluricanonical differential forms on $\overline{\mathcal{F}}_L(\Gamma)$. Any such differential form can be obtained using a modular form (see the proof of Theorem 6.13). The differential form $F_{n,m}(dZ)^m$ is Γ -invariant and it determines a section of the pluricanonical bundle mK over a smooth open

part of the modular variety away from the branch locus of $\pi: \mathcal{D}_L \rightarrow \mathcal{F}_L(\Gamma)$ and the cusps. In the proof of Theorem 6.13 we indicated three kinds of obstruction to extending $F_{n,m} (dZ)^m$ to a global section of mK . In the proof of this theorem we use the reflective obstruction, arising from the ramification divisor in \mathcal{D}_L by \pm reflections in Γ (see Equation (6.12)). Therefore if $F_{n,m}$ determines a global section then $F_{n,m}$ has zeroes of order at least m on $\text{Bdiv}(\pi_\Gamma)$. The modular form $F_k \in M_k(\Gamma, \chi)$ is strongly reflective of weight $k \geq n$. Hence $F_{n,m}/F_k^m$ is a holomorphic modular form of weight $m(n - k) \leq 0$. According to Koecher’s principle ($n \geq 3$) this function is constant. Therefore $F_{n,m} \equiv 0$ if $k > n$ or $F_{n,m} = C \cdot F_n^m$ if $k = n$. If the strongly reflective form F_n is non-cuspidal of weight n , then $F_n^m (dZ)^{\otimes m}$ cannot be extended to the compact model because of cusp obstructions (F_n^m should have zeroes of order at least m along the boundary). If F_n is a cusp form of weight k then we can consider F_n as a cusp form with respect to the subgroup Γ_χ .

Then $F_n(Z) dZ$ is Γ_χ -invariant and, according to Freitag’s lemma, it can be extended to a global section of the canonical bundle $\Omega_{\overline{\mathcal{F}}_L(\Gamma_\chi)}$ for any smooth compact model $\overline{\mathcal{F}}_L(\Gamma_\chi)$ of $\mathcal{F}_L(\Gamma_\chi)$. Moreover Koecher’s principle shows that any m -pluricanonical form is equal, up to a constant, to $F_n^m (dZ)^{\otimes m}$, proving (6.21). The strongly reflective cusp form of canonical weight determines essentially the unique m -pluricanonical differential form on $\mathcal{F}_L(\Gamma_\chi)$. □

We can apply Theorem 6.19 to find examples of moduli spaces of lattice-polarised K3 surfaces having $\kappa = -\infty$ and $\kappa = 0$: see [41]. Meanwhile a stronger statement concerning uniruledness of orthogonal varieties has been proven, see [45].

7. Orthogonal groups and reflections

The material in this and subsequent sections is not so easily found in the literature, so from here on we shall give slightly more detail.

For applications, the most important subgroups of $O(L)$ are the stable orthogonal groups $\tilde{O}(L)$, $\tilde{O}^+(L)$ and $\tilde{SO}^+(L)$, as defined in Equations (2.14) and (2.15). The reason for using the word “stable” to describe $\tilde{O}(L)$ is the following property.

Lemma 7.1. *For any sublattice S of a lattice L the group $\tilde{O}(S)$ can be considered as a subgroup of $\tilde{O}(L)$.*

Proof. Let S^\perp be the orthogonal complement of S in L . We have

$$(7.2) \quad S \oplus S^\perp \subset L \subset L^\vee \subset S^\vee \oplus (S^\perp)^\vee$$

where $S \oplus S^\perp$ is a sublattice of finite index in L . We can extend $g \in \tilde{O}(S)$ on $S \oplus S^\perp$ putting $g|_{S^\perp} \equiv \text{id}$. It is clear that $g \in \tilde{O}(S \oplus S^\perp)$. We consider $g \in \tilde{O}(S \oplus S^\perp)$ as an element of $O(S^\vee \oplus (S^\perp)^\vee)$. For any $l^\vee \in L^\vee$ we have $g(l^\vee) \in l^\vee + (S \oplus S^\perp)$. In particular, $g(l) \in L$ for any $l \in L$ and $g \in \tilde{O}(L)$. □

Let S be a primitive sublattice of L . We define the groups

$$O(L, S) = \{g \in O(L) \mid g|_S \in \tilde{O}(S)\} \quad \text{and} \quad \tilde{O}(L, S) = O(L, S) \cap \tilde{O}(L).$$

Note that $O(L, \mathbb{Z}h) = O(L, h)$ if $h^2 \neq \pm 2$. The technique of discriminant forms developed by Nikulin in [97] is very useful here, and we describe the main ideas behind it below. For simplicity we assume that all the lattices we consider are even.

Let S^\perp be the orthogonal complement of a primitive nondegenerate sublattice S in L . As in the proof of Lemma 7.1 we have the inclusions (7.2). The overlattice L is defined by the finite subgroup

$$(7.3) \quad H = L/(S^\perp \oplus S) < (S^\perp)^\vee/S^\perp \oplus S^\vee/S = D(S^\perp) \oplus D(S)$$

which is an isotropic subgroup of $D(S^\perp) \oplus D(S)$. Moreover $L/(S \oplus S^\perp) \cong L^\vee/(S^\vee \oplus (S^\perp)^\vee)$. We define $\phi: L \rightarrow S^\vee$ by $\phi(l)(s) = (l, s)$. Then $\ker(\phi) = S^\perp$. Since $L/(S \oplus S^\perp) \cong \phi(L)/S$ we obtain

$$|L/(S \oplus S^\perp)| = |\phi(L)/S| = |\det S|/[S^\vee : \phi(L)],$$

as $|\det S| = [S^\vee : S]$. From the inclusions above

$$|\det S| \cdot |\det S^\perp| = (|\det L|)[\phi(L) : S]^2 = |\det L| \cdot |\det S|^2/[S^\vee : \phi(L)]^2.$$

In the particular case $S = \mathbb{Z}h$ and $L_h = h^\perp$ we have $[S^\vee : \phi(L)] = \text{div}(h)$, where

$$(7.4) \quad \text{div}(h)\mathbb{Z} = (h, L).$$

The positive number $\text{div}(h)$ is called the *divisor* of h in L . We have now proved the following lemma.

Lemma 7.5. *Let L be any nondegenerate even integral lattice and let $h \in L$ be a primitive vector with $h^2 = 2d \neq 0$. If L_h is the orthogonal complement of h in L then*

$$|\det L_h| = \frac{|(2d) \cdot \det L|}{\text{div}(h)^2}.$$

We come back to the inclusion (7.3). Following [97] we consider the projections

$$p_S: H \rightarrow D(S), \quad p_{S^\perp}: H \rightarrow D(S^\perp).$$

Using the definitions and the fact that the lattices S and S^\perp are primitive in L one can show (see [97, Prop. 1.5.1]) that these projections are injective and moreover that if $d_S \in p_S(H)$ then there is a unique $d_{S^\perp} \in p_{S^\perp}(H)$ such that $d_S + d_{S^\perp} \in H$. Using these arguments one proves the next lemma (see [50, Lemma 3.2])

Lemma 7.6. *Let S be a primitive sublattice of an even lattice L and denote by \bar{g} the image in $O(D(L))$ of $g \in O(L)$.*

- (i) $g \in O(L, S)$ if and only if $g(S) = S$, $\bar{g}|_{D(S)} = \text{id}$ and $\bar{g}|_{p_{S^\perp}(H)} = \text{id}$.
- (ii) $\alpha \in O(S^\perp)$ can be extended to $O(L, S)$ if and only if $\bar{\alpha}|_{p_{S^\perp}(H)} = \text{id}$.
- (iii) If $p_{S^\perp}(H) = D(S^\perp)$ then $O(L, S)|_{S^\perp} \cong \tilde{O}(S^\perp)$.

(iv) Assume that the projection $O(S^\perp) \rightarrow O(D(S^\perp))$ is surjective. Then

$$O(L, S)|_{S^\perp} / \widetilde{O}(S^\perp) \cong \{\overline{\gamma} \in O(D(S^\perp)) \mid \overline{\gamma}|_{\mathfrak{p}_{S^\perp}(H)} = \text{id}\}.$$

Corollary 7.7. *If $|H| = |\det S^\perp|$ then $O(L, S)|_{S^\perp} \cong \widetilde{O}(S^\perp)$.*

For example, the condition of Corollary 7.7 is true if L is an even unimodular lattice and S is any primitive sublattice of L (see Example 7.9 below). We recall the following result which we call *Eichler’s criterion* (see [30, Section 10], [40, Section 3] and [49]).

Lemma 7.8. *Let L be a lattice containing two orthogonal hyperbolic planes. Then the $\widetilde{SO}(L)$ -orbit of a primitive vector $l \in L$ is determined by two invariants: by its length $l^2 = (l, l)$ and its image $l^* + L$, where $l^* = l / \text{div}(l)$, in the discriminant group $D(L)$.*

We note that l^* is a primitive element of the dual lattice L^\vee . Therefore $\text{div}(l)$ is a divisor of the exponent of the discriminant group $D(L)$. In particular, $\text{div}(l)$ divides $\det(L)$. Lemma 7.8 can be used to classify all possible vectors of fixed length in different lattices.

Example 7.9. (*The K3 lattice.*) $L(K3) = 3U \oplus 2E_8(-1)$ is the even unimodular lattice of signature $(3, 19)$: its discriminant group is trivial and all the primitive vectors $h_{2d} \in L_{K3}$ of length $2d$ form a single $SO(L_{K3})$ -orbit. Therefore we can take h_{2d} in the first hyperbolic plane U , so

$$(7.10) \quad (h_{2d})_{L_{K3}}^\perp \cong L_{2d} = 2U \oplus 2E_8(-1) \oplus \langle -2d \rangle.$$

Then according to Corollary 7.7

$$(7.11) \quad O^+(L_{K3}, h_{2d}) \cong \widetilde{O}^+(h_{2d}^\perp) = \widetilde{O}^+(2U \oplus 2E_8(-1) \oplus \langle -2d \rangle).$$

The case of K3 surfaces is an exception, since the K3 lattice is unimodular.

Example 7.12. $K3^{[2]}$ -lattice, split and non-split polarisations. We consider the Beauville lattice of a deformation $K3^{[n]}$ manifold, which is isomorphic to $L_{K3, 2n-2}$ by Proposition 3.2(i). In general there will be several, but finitely many, orbits of primitive polarisation vectors h_{2d} . Let $h_{2d} \in L_{K3, 2}$ be a primitive vector of length $2d > 0$. Then $\text{div}(h_{2d})$ divides $|\det L_{K3, 2}| = 2$.

All vectors with $\text{div}(h_{2d}) = 1$ constitute a single $\widetilde{SO}^+(L_{K3, 2})$ -orbit, by Lemma 7.8. Therefore, as in Example 7.9, we obtain

$$(7.13) \quad (h_{2d})_{L_{K3, 2}}^\perp \cong L_{2, 2d} = 2U \oplus 2E_8(-1) \oplus \langle -2 \rangle \oplus \langle -2d \rangle.$$

We call a polarisation determined by a primitive vector h_{2d} with $\text{div}(h_{2d}) = 1$ a *split polarisation*.

If $h_{2d} \in L_{K3, 2}$ and $\text{div}(h_{2d}) = 2$ then we can write h_{2d} as $h_{2d} = 2v + cl_2$, where $v \in 3U \oplus 2E_8(-1)$ and l_2 is a generator of the orthogonal component $\langle -2 \rangle$ in $L_{K3, 2}$. The coefficient c is odd because h_{2d} is primitive. Note that $2d = h_{2d}^2 = 4(v, v) - 2c^2$, so $d \equiv -1 \pmod{4}$. According to Eichler’s criterion the $\widetilde{SO}(L_{K3, 2})$ -orbit of h_{2d} is

uniquely determined by the class $h_d^* \equiv l_2/2 \pmod{L_{K3,2}}$. Therefore as in the case of $\text{div}(h_{2d}) = 1$ all vectors with $\text{div}(h_{2d}) = 2$ form only one orbit. We can take a representative in the form $h_{2d} = 2v + cl_2 \in U \oplus \langle -2 \rangle$. The orthogonal complement of h_{2d} in $U \oplus \langle -2 \rangle$ can be found by direct calculation. This is an even rank 2 lattice $Q(d)$ of determinant d . It follows that if $d \equiv -1 \pmod{4}$ then there is only one orbit of vectors h_{2d} with $\text{div}(h_{2d}) = 2$, and the orthogonal complement of h_{2d} is uniquely determined:

$$(7.14) \quad (h_{2d})_{L_{K3,2}}^\perp \cong L_{Q(d)} = 2U \oplus 2E_8(-1) \oplus \begin{pmatrix} -2 & 1 \\ 1 & -\frac{d+1}{2} \end{pmatrix}.$$

We call a polarisation of this kind a *non-split polarisation*. We note that $|\det L_{Q(d)}| = d$ and the discriminant group of $L_{Q(d)}$ is cyclic (see [50, Remark 3.15]).

Remark 7.15. Taking the orthogonal complement of the (-2) -vector in $Q(d)$ we have a split sublattice of index 2

$$\langle -2 \rangle \oplus \langle -2d \rangle < Q(d).$$

Therefore $L_{2,2d} < L_{Q(d)}$ is also a sublattice of index 2 and according to Lemma 7.1

$$(7.16) \quad \tilde{O}^+(L_{2,2d}) < \tilde{O}^+(L_{Q(d)})$$

is a subgroup of finite index. This implies that the modular variety $\mathcal{F}_L(\tilde{O}^+(L_{2,2d}))$ is a finite covering of $\mathcal{F}_L(\tilde{O}^+(L_{Q(d)}))$ where $d \equiv -1 \pmod{4}$. One can calculate this index using the explicit formula for the Hirzebruch–Mumford volume (see [47]).

We gave a classification of all possible type of polarisations for the symplectic varieties of $K3^{[n]}$ type in [50, Proposition 3.6]. Results about the polarisation types of 10-dimensional O’Grady varieties can be found in [51, Theorem 3.1].

The branch divisor (Equation (6.12)) of the modular varieties is defined by \pm reflections in the modular groups. Below we give a description of the branch divisors in the cases of the polarised K3 surfaces and polarised holomorphic symplectic varieties of type $K3^{[2]}$.

Let L be a nondegenerate integral lattice and $r \in L$ be a primitive vector. If the reflection is integral, i.e. $\sigma_r \in O(L)$ (see Equations (6.11) and (7.4)), then

$$(7.17) \quad \text{div}(r) \mid r^2 \mid 2 \text{div}(r).$$

The following general result was proved in [46].

Proposition 7.18. (i) *Let L be a nondegenerate even integral lattice. Let $r \in L$ be primitive. Then $\sigma_r \in \tilde{O}(L)$ if and only if $r^2 = \pm 2$.*

If $-\sigma_r \in \tilde{O}(L)$, i.e. $\sigma_r|_{A_L} = -\text{id}$, then we also have

- (ii) $r^2 = \pm 2a$ and $\text{div}(r) = a \equiv 1 \pmod{2}$, or $r^2 = \pm a$ and $\text{div}(r) = a$ or $a/2$; and
- (iii) $A_L \cong (\mathbb{Z}/2\mathbb{Z})^m \times (\mathbb{Z}/a\mathbb{Z})$, for some $m \geq 0$.

If (iii) holds then

- (iv) If $r^2 = \pm a$ and either $\text{div}(r) = a$ or $\text{div}(r) = a/2 \equiv 1 \pmod 2$, then $-\sigma_r \in \tilde{O}(L)$;
- (v) If $r^2 = \pm 2a$ and $\text{div}(r) = a \equiv 1 \pmod 2$, then $-\sigma_r \in \tilde{O}(L)$.

With polarised K3 surfaces in mind, we consider in more detail the lattice L_{2d} (see Equation (7.10)). In this case the ramification divisor has three irreducible components.

Corollary 7.19. *Let σ_r be a reflection in $O^+(L_{2d})$ defined by a primitive vector $r \in L_{2d}$. The reflection σ_r induces $\pm \text{id}$ on the discriminant group L_{2d}^\vee/L_{2d} if and only if $r^2 = -2$ or $r^2 = -2d$ and $\text{div}(r) = d$ or $2d$. If $r^2 = -2$ then*

$$r_{L_{2d}}^\perp \cong 2U \oplus E_8(-1) \oplus E_7(-1) \oplus \langle -2d \rangle.$$

If $\text{div}(r) = d$ then either

$$r_{L_{2d}}^\perp \cong U \oplus 2E_8(-1) \oplus \langle 2 \rangle \oplus \langle -2 \rangle$$

or

$$r_{L_{2d}}^\perp \cong U \oplus 2E_8(-1) \oplus U(2).$$

See the proof in [46, Corollary 3.4 and Proposition 3.6]. Geometrically the three cases in the last proposition correspond to the Néron-Severi group being (generically) U , $U(2)$ or $\langle 2 \rangle \oplus \langle -2 \rangle$ respectively. The K3 surfaces (without polarisation) themselves are, respectively, a double cover of the Hirzebruch surface F_4 , a double cover of a quadric, and the desingularisation of a double cover of \mathbb{P}^2 branched along a nodal sextic.

We note that in the case of polarised deformation K3^[2] manifolds the branch divisor has one main (i.e. $\text{div}(r) = 1$) component, with $r^2 = -2$, and 6 (respectively, 1) additional components ($\text{div}(r) > 1$) for split (respectively, non-split) type (see the proof of Proposition 8.22 below).

8. The quasi pull-back of modular forms

The main aim of this section is to show how we can construct cusp forms of small weight, for example on the moduli spaces of polarised K3 surfaces. We use the method of quasi pull-back of the Borcherds form Φ_{12} which was proposed in [11, pp. 200-201]. This method was successfully applied to the theory of moduli spaces in [14], [76], [46], [50] and [51]. In this section we review this method and prove a new result (Theorem 8.18 below) showing that non-trivial quasi pull-backs are cusp forms.

8.1. Quasi pull-back

First we give a general property of rational quadratic divisors. Let M be a lattice of signature $(2, m)$ and L be a primitive nondegenerate sublattice of signature

$(2, n)$ where $n < m$. Then L_M^\perp is negative definite and we have the usual inclusions $L \oplus L_M^\perp < M < M^\vee < L^\vee \oplus (L^\perp)^\vee$. For $v \in M$ we write

$$(8.1) \quad v = \alpha + \beta, \quad \alpha = \text{pr}_{L^\vee}(v) \in L^\vee, \beta \in (L^\perp)^\vee.$$

Lemma 8.2. *Let L and M be as above. Then for any $v \in M$ with $v^2 < 0$ we have*

$$\mathcal{D}_L \cap \mathcal{D}_v(M) = \begin{cases} \mathcal{D}_\alpha(L), & \text{if } \alpha^2 < 0, \\ \emptyset, & \text{if } \alpha^2 \geq 0, \alpha \neq 0, \\ \mathcal{D}_L, & \text{if } \alpha = 0, \text{ i.e. } v \in L^\perp. \end{cases}$$

Proof. We have $\mathcal{D}_L \subset \mathcal{D}_M$ because L is a sublattice of M . For any $Z = X + iY \in \mathcal{D}_L^\bullet$ with $X, Y \in L \otimes \mathbb{R}$ we have $\langle X, Y \rangle = 0$ and $\langle X, X \rangle = \langle Y, Y \rangle > 0$. Therefore the quadratic space $\langle X, Y \rangle_{\mathbb{R}}$ is of signature $(2, 0)$. Note that $\langle Z, v \rangle = 0$ is equivalent to $\langle Z, \alpha \rangle = 0$. The signature of L is equal to $(2, n)$. Analysing the signature of $\langle X, Y \rangle_{\mathbb{R}} \oplus \langle \alpha \rangle_{\mathbb{R}}$ we get that $\mathcal{D}_L \cap \mathcal{D}_v(M)$ is non-empty if and only if $\alpha = \text{pr}_{L^\vee}(v)$ belongs to the negative definite quadratic space $\langle X, Y \rangle_{M \otimes \mathbb{R}}^\perp$. This proves the first two cases of the lemma.

The finite group $H = M/(L \oplus L^\perp)$ is a subgroup of the orthogonal sum of the discriminant groups $D(L) \oplus D(L^\perp)$ where $D(L) = L^\vee/L$. The decomposition (8.1) defines a projection $\text{pr}: H \rightarrow D(L) \oplus D(L^\perp)$. For a primitive sublattice L the class $(\alpha + L) \in D(L)$ is uniquely determined by the class $\beta + L^\perp$ in $D(L^\perp)$ (see [97, Proposition 1.5.1]). Therefore $\alpha \in L$ if and only if $\beta \in L^\perp$. In particular v is orthogonal to L if and only if $\alpha = 0$. This proves the last assertion of the lemma. \square

In the next theorem we explain the main idea of the method of quasi pull-back applied to the strongly reflective modular form Φ_{12} (see [11, pp. 200–201] and [14]).

Theorem 8.3. *Let $L \hookrightarrow \text{II}_{2,26}$ be a primitive nondegenerate sublattice of signature $(2, n)$, $n \geq 3$, and let $\mathcal{D}_L \hookrightarrow \mathcal{D}_{\text{II}_{2,26}}$ be the corresponding embedding of the homogeneous domains. The set of (-2) -roots*

$$R_{-2}(L^\perp) = \{r \in \text{II}_{2,26} \mid r^2 = -2, (r, L) = 0\}$$

in the orthogonal complement is finite. We put $N(L^\perp) = \#R_{-2}(L^\perp)/2$. Then the function

$$(8.4) \quad \Phi|_L = \frac{\Phi_{12}(Z)}{\prod_{r \in R_{-2}(L^\perp)/\pm 1} (Z, r)} \Big|_{\mathcal{D}_L} \in M_{12+N(L^\perp)}(\tilde{\mathcal{O}}(L), \det),$$

where in the product over r we fix a finite system of representatives in $R_{-2}(L^\perp)/\pm 1$. The modular form $\Phi|_L$ vanishes only on rational quadratic divisors of type $\mathcal{D}_v(L)$ where $v \in L^\vee$ is the orthogonal projection of a (-2) -root $r \in \text{II}_{2,26}$ on L^\vee .

We say that the modular form $\Phi|_L$ is a *quasi pull-back* of Φ_{12} if the set of roots $R_{-2}(L^\perp)$ is non-empty.

Proof. We introduce coordinates $Z = (Z_1, Z_2) \in \mathcal{D}_{\text{II}_{2,26}}$ related to the embedding $L \hookrightarrow \text{II}_{2,26}$ and the splitting (8.1), namely $Z_1 \in L \otimes \mathbb{C}$ and $Z_2 \in (L \otimes \mathbb{C})^\perp$. We have $\tilde{\mathcal{O}}^+(L) < \mathcal{O}^+(\text{II}_{2,26})$ (see Lemma 7.1) and we denote by $\tilde{g} \in \mathcal{O}^+(\text{II}_{2,26})$ the extension of $g \in \tilde{\mathcal{O}}^+(L)$ by $\tilde{g}|_{L^\perp} = \text{id}$.

If the root system $R_{-2}(L^\perp)$ is empty, then $\Phi|_L$ is the usual pull-back of Φ_{12} on \mathcal{D}_L . Then $\Phi|_L \neq 0$ and we have

$$\det(g)\Phi_{12}(Z_1, Z_2) = \Phi_{12}(\tilde{g}(Z_1, Z_2)) = \Phi_{12}(g \cdot Z_1, Z_2).$$

Therefore the pull-back of Φ_{12} on \mathcal{D}_L is a modular form of weight 12. We note that even in this simple case one obtains interesting reflective modular forms (see [55, Section 4.2]).

If there are roots in $R_{-2}(L^\perp)$ then the pull-back of Φ_{12} on \mathcal{D}_L vanishes identically, and one has to divide by the equations of the rational quadratic divisors, as in Equation (8.4).

According to Lemma 8.2 the order of zero of Φ_{12} along \mathcal{D}_L is equal to $N(L^\perp) = \#R_{-2}(L^\perp)/2$. Therefore the non-zero function $\Phi|_L$ is holomorphic on \mathcal{D}_L^\bullet . Moreover it is homogeneous of degree $12 + N(L^\perp)$. Any $g \in \tilde{\mathcal{O}}^+(L)$ acts trivially on L^\perp and $(Z, r) = (Z_1 + Z_2, r) = (Z_2, r)$. Therefore

$$\left. \frac{\Phi_{12}(gZ_1, Z_2)}{\prod_r(Z_2, r)} \right|_{\mathcal{D}_L} = \left. \frac{\Phi_{12}(\tilde{g} \cdot (Z_1, Z_2))}{\prod_r(\tilde{g} \cdot (Z_1, Z_2), r)} \right|_{\mathcal{D}_L} = \det(g) \left. \frac{\Phi_{12}(Z)}{\prod_r(Z, r)} \right|_{\mathcal{D}_L}.$$

It follows that $\Phi|_L$ is modular with respect to $\tilde{\mathcal{O}}^+(L)$ with character \det . We finish the proof using Koecher’s principle.

The zeros of $\Phi|_L$ can be determined using Lemma 8.2 and the fact that Φ_{12} vanishes along $\mathcal{D}_r(\text{II}_{2,26})$ with $r^2 = -2$. □

Remark 8.5. Theorem 8.3 is still true for $n \leq 2$. We can show this by computing the Fourier expansions of the quasi pull-back. Moreover we prove in Theorem 8.18 that the quasi pull-back is always a cusp form.

Remark 8.6. The modular group of $\Phi|_L$ might be larger than $\tilde{\mathcal{O}}^+(L)$ (see, for example, [51, Lemma 4.4]).

Remark 8.7. For the applications to the theory of moduli spaces we use the quasi pull-back of Φ_{12} . It is easy to prove an analogue of Theorem 8.3 for an arbitrary modular form whose divisor consists only of rational quadratic divisors (in the style of Theorem 8.18 below).

In [46] we showed that some quasi pull-backs for lattices related to the moduli spaces of polarised K3 surfaces are cusp forms. In this paper we prove a new, more general result: that the quasi pull-back construction always gives a cusp form. The main idea of the proof is to consider the quasi pull-back as a differential operator (see [46, Section 6]).

8.2. Tube domain realisation

We define a Fourier expansion of a modular form F at a 0-dimensional cusp. The Fourier expansion depends on the choice of affine coordinates at a cusp. We consider the general case, following the approach used in [12, Theorem 5.2 and page 542]: for more details see [55, Section 2.3] and [33].

A 0-dimensional cusp of \mathcal{D}_L is defined by an isotropic sublattice of rank 1 or, equivalently, by a primitive isotropic vector $c \in L$ (up to sign: c and $-c$ define the same cusp). The choice of c identifies \mathcal{D}_L with an affine quadric:

$$\mathcal{D}_{L,c} = \{Z \in \mathcal{D}_L^\bullet \mid (Z, c) = 1\} \cong \mathcal{D}_L.$$

The lattice

$$(8.8) \quad L_c := c^\perp / c = c_L^\perp / \mathbb{Z}c$$

is an integral lattice of signature $(1, n-1)$. We fix an element $b \in L^\vee$ such that $(c, b) = 1$. A choice of b gives a realisation of the hyperbolic lattice L_c as a sublattice in L

$$(8.9) \quad L_c \cong L_{c,b} = L \cap c^\perp \cap b^\perp.$$

We have

$$L \otimes \mathbb{Q} = L_{c,b} \otimes \mathbb{Q} \oplus (\mathbb{Q}b + \mathbb{Q}c).$$

Using the hyperbolic lattice $L_c \otimes \mathbb{R}$ we define a positive cone

$$C(L_c) = \{x \in L_c \otimes \mathbb{R} \mid (x, x) > 0\}.$$

We may choose $C^+(L_c)$, one of the two connected components of $C(L_c)$ so that corresponding tube domain, which is the complexification of $C^+(L_c)$

$$\mathcal{H}_c = L_c \otimes \mathbb{R} + iC^+(L_c)$$

has an isomorphism $\mathcal{H}_c \rightarrow \mathcal{D}_{L,c} \cong \mathcal{D}_L$ by

$$(8.10) \quad z \mapsto [z] = z \oplus \left(b - \frac{(z, z) + (b, b)}{2} c \right) \quad (z \in \mathcal{H}_c, [z] \in \mathcal{D}_{L,c}).$$

Using the coordinate $z \in \mathcal{H}_c$ defined by the choice of c and b we can identify an arbitrary modular form F of weight k with a modular form $F_{c,b}$ (or simply F_c) on the tube domain \mathcal{H}_c :

$$F(\lambda[z]) = \lambda^{-k} F_{c,b}(z).$$

8.3. Fourier expansion at 0-dimensional cusps

In order to define the Fourier expansion at the cusp c we consider an unipotent subgroup of the stabiliser $\tilde{O}^+(L)$, the subgroup of the Eichler transvections. For any $a \in c_L^\perp$ the map

$$t'(c, a): v \mapsto v - (a, v)c \quad (v \in c_L^\perp)$$

belongs to the orthogonal group $O(c_\perp^\perp)$. It has the unique orthogonal extension on L which is given by the map

$$(8.11) \quad t(c, a): v \mapsto v - (a, v)c + (c, v)a - \frac{1}{2}(a, a)(c, v)c.$$

This element is called an *Eichler transvection*: see [30, Section 3] and [49]. We note that $t(c, a) \in \widetilde{SO}^+(L)$ for any $a \in c_\perp^\perp$, that $t(c, a)(c) = c$, and that for $a, a' \in c_\perp^\perp$

$$t(c, a)t(c, a') = t(c, a + a') \quad \text{and} \quad t(c, a)^{-1} = t(c, -a).$$

We can identify the lattice $L_{c,b}$ with the corresponding group of transvections $E_c(L) = \langle t(c, a) \mid a \in L_c \rangle$. The group $E_c(L)$ is the unipotent radical of the parabolic subgroup associated to c . A direct calculation shows that $t(c, a)$ acts as linear translation in the affine coordinates (8.10):

$$t(c, a)([z]) = [z + a].$$

Let $F \in M_k(\widetilde{SO}^+(L))$. Then $F_c(z + a) = F_c(z)$ for all $a \in L_{c,b}$ and we obtain the Fourier expansion of F at the cusp c :

$$(8.12) \quad F_c(z) = \sum_{l \in L_{c,b}^\vee} f(l) \exp(2\pi i(l, z)).$$

The function $F_c(z)$ is holomorphic at the cusp c if the Fourier coefficient $f(l)$ can be different from 0 only if its index $l \in L_{c,b}^\vee$ belongs to the closure of the positive cone $C^+(L_c)$. Another formulation of this fact is $(l, Y) > 0$ for any Y in the positive cone $C^+(L_c)$.

Remark 8.13. In general the Fourier expansion (8.12) depends on the choice of b . For another b' the Fourier coefficients will be different by a factor $\exp(2\pi i(l, b - b'))$ which is a root of unity (see details in [55, §2.3]). In particular the Fourier coefficient f_0 (the value of F at the cusp c) is well-defined.

Remark 8.14. The stabiliser of the cusp c in $\widetilde{O}^+(L)$ is isomorphic to the semi-direct product of $\widetilde{O}^+(L_c)$ and $E_c(L)$.

Remark 8.15. Let $\text{div}_L(c) = N$. Then $\det L = N^2 \det L_c$. If $\text{div}_L(c) = 1$ then c can be completed to a hyperbolic plane in L and $L = U \oplus L_c$. This cusp is called the *simplest cusp*. If $\text{div}(c) > 1$ then $c/\text{div}(c) + L$ is an isotropic element of the discriminant group $D(L)$ and $|D(L)| = |\det L|$ is divisible by $\text{div}(c)^2$. If $D(L)$ contains no non-trivial isotropic elements (in this case the lattice is maximal) then all 0-dimensional cusps are equivalent to the simplest cusp.

Remark 8.16. If L contains two hyperbolic planes then one can use the Eichler criterion (see Lemma 7.8) in order to classify the 0-dimensional cusps with respect to the action of $\widetilde{SO}^+(L)$. The orbit of c , in this case, is uniquely determined by the isotropic class $c/\text{div}(c) + L$ in the discriminant group $D(L)$.

If $\mathcal{D}(L)$ does not contain isotropic elements (in particular if $\det(L)$ is square free) then the modular variety $\widetilde{\mathcal{O}}^+(L) \setminus \mathcal{D}_L$ has only one 0-dimensional cusp.

Remark 8.17. If $[\mathcal{O}(L) : \Gamma] < \infty$ and $F \in M_k(\Gamma, \chi)$ then we can define the Fourier expansion using a sublattice $L_c(\Gamma)$ of finite index in L_c corresponding to the group $E_c(L) \cap \ker \chi$.

8.4. Properties of quasi pull-back

We may consider the quasi pull-back of other modular forms, not only Φ_{12} . If L is of signature $(2, n)$ and $r \in L$ is primitive with $(r, r) < 0$ then $\mathcal{D}_r(L) = \mathcal{D}_{tr}(L)$ for any $t \in \mathbb{Q}^\times$, and we write $r^\perp = r^\perp_1$ if there is no ambiguity.

Theorem 8.18. *Let L be an integral lattice of signature $(2, n)$. Suppose that the modular form $F \in M_k(\widetilde{\mathcal{S}\mathcal{O}}^+(L))$ vanishes with order $m > 0$ on the rational quadratic divisor $\mathcal{D}_r(L)$ where r is a primitive vector in L with $(r, r) < 0$. We define the quasi pull-back of F on the domain \mathcal{D}_{r^\perp} of complex dimension $n - 1$ by*

$$(8.19) \quad F|_{r^\perp} = \frac{F(Z)}{(Z, r)^m} \Big|_{\mathcal{D}_{r^\perp}}.$$

Then the quasi pull-back is a cusp form of weight $k + m$

$$F|_{r^\perp} \in S_{k+m}(\widetilde{\mathcal{S}\mathcal{O}}^+(r^\perp)).$$

Proof. There are two parts to the assertion: that $F|_{r^\perp}$ is a modular form, and that it vanishes at every cusp.

For the first part, we have the inclusions

$$r^\perp \oplus \mathbb{Z}r \subset L \subset L^\vee \subset (r^\perp)^\vee \oplus \mathbb{Z} \frac{r}{(r, r)},$$

and r^\perp has signature $(2, n - 1)$. We consider the two embeddings $\mathcal{D}_{r^\perp} \hookrightarrow \mathcal{D}_L$ and $\widetilde{\mathcal{S}\mathcal{O}}^+(r^\perp) \hookrightarrow \widetilde{\mathcal{S}\mathcal{O}}^+(L)$. We note that any element $g \in \widetilde{\mathcal{S}\mathcal{O}}^+(r^\perp)$ extends to $\tilde{g} \in \widetilde{\mathcal{S}\mathcal{O}}^+(L)$ by acting trivially on r (see Lemma 7.1). Therefore any \tilde{g} preserves \mathcal{D}_{r^\perp} and $\tilde{g} \cdot r = r$. The function $F|_{r^\perp}$ is a holomorphic function on \mathcal{D}_{r^\perp} and it is homogeneous of degree $k + m$. For any $g \in \widetilde{\mathcal{S}\mathcal{O}}^+(r^\perp)$ we have

$$\frac{F(\tilde{g}Z)}{(\tilde{g}Z, r)^m} \Big|_{\mathcal{D}_{r^\perp}} = \frac{F(Z)}{(Z, r)^m} \Big|_{\mathcal{D}_{r^\perp}}.$$

Therefore the quasi pull-back $F|_{r^\perp}$ is $\widetilde{\mathcal{S}\mathcal{O}}^+(r^\perp)$ -invariant. If instead F has character χ then $F|_{r^\perp}$ transforms according to the induced character $\chi|_{\widetilde{\mathcal{S}\mathcal{O}}^+(r^\perp)}$. If $n > 3$ then using Koecher's principle we conclude that

$$F|_{r^\perp} \in M_{k+m}(\widetilde{\mathcal{S}\mathcal{O}}(r^\perp)).$$

If $n \leq 3$ we cannot apply Koecher's principle and instead we must use the Fourier expansion to check that the quasi pull-back is a modular form as well as to show the vanishing at the cusps.

We calculate the Fourier expansion of the quasi pull-back at an arbitrary 0-dimensional cusp of r^\perp . Let $c \in r^\perp$ be a primitive isotropic vector in r^\perp (if there are any: if not, there is nothing more to prove). We fix a vector $b \in (r^\perp)^\vee$. Then we can define the two homogeneous domains $\mathcal{H}_c(r^\perp)$ and $\mathcal{H}_c(L)$ because the vector c defines also a 0-dimensional cusp of L . We write $Z \in \mathcal{H}_c(L)$ in the form $Z = Z_1 + zr$ where

$$Z_1 \in r^\perp \otimes \mathbb{C}, \quad z = x + iy \in \mathbb{C}, \quad (\text{Im } Z_1, \text{Im } Z_1) + (r, r)y^2 > 0.$$

If $[Z]$ is the image of Z in $\mathcal{D}_c(L)$ (see (8.10)) then $([Z], r) = (r, r)z$. Therefore the equation of the divisor $\mathcal{D}_r(L)$ in the affine coordinates $Z = Z_1 \oplus zr \in \mathcal{H}_c$ is $z = 0$. The quasi pull-back $F|_{r^\perp}$ is equal, up to a constant, to the first non-zero coefficient in the Taylor expansion of $F_{c,b}(Z)$ in z .

Consider the Fourier expansion

$$F_{c,b}(Z) = \sum_{l \in L_{c,b}^\vee} f(l) \exp(2\pi i (l, Z)).$$

The modular form $F_{c,b}(Z)$ is holomorphic at the boundary. Therefore l belongs to the closure of the dual cone, in particular, $(l, l) \geq 0$. The vectors c and b are orthogonal to r and the lattice $(r^\perp)_{c,b} \oplus \mathbb{Z}r$ is a sublattice of finite index in the lattice of translations $L_{c,b}$. For any $Z_1 \in \mathcal{H}_c(r^\perp) \subset r^\perp \otimes \mathbb{C}$ we consider $z = x + iy$ such that $Z = Z_1 \oplus zr \in \mathcal{H}_c(L)$. (If y is small enough then $(\text{Im } Z_1, \text{Im } Z_1) > -(r, r)y^2 > 0$.) Therefore we can rewrite the Fourier expansion using this parametrisation

$$(8.20) \quad F_{c,b}(Z_1 + zr) = \sum_{l_1 \in (r^\perp)_{c,b}^\vee, l_2 \in \mathbb{Z}r/(r,r)} f(l_1 + l_2) \exp(2\pi i (l_1, Z_1) + z(l_2, r))$$

where $(l, l) = (l_1 + l_2, l_1 + l_2) = (l_1, l_1) + (l_2, l_2) \geq 0$. The Fourier coefficients of $(F|_{r^\perp})_{c,b}$ are proportional to the Fourier coefficients of the m -th Taylor coefficient

$$\left. \frac{\partial^m F_{c,b}(Z_1 + zr)}{(\partial z)^m} \right|_{z=0}.$$

The derivatives of the terms in the Fourier expansion (8.20) vanish if $l_2 = 0$. If $(l_2, l_2) < 0$ then $(l_1, l_1) > 0$, so nonzero Fourier coefficients occur only when the index l_1 has positive square. We have proved this, which is much stronger than just the vanishing of the zeroth coefficient, for an arbitrary 0-dimensional cusp of r^\perp , so we have shown that $F|_{r^\perp}$ is holomorphic at the boundary even for $n = 1$ or $n = 2$. Moreover the value of $F|_{r^\perp}$ at an arbitrary 0-dimensional cusp is zero.

The Baily-Borel compactification of $\widetilde{SO}^+(r^\perp) \setminus \mathcal{D}_{r^\perp}$ contains only boundary components of dimension 0 and 1 (the latter only if r^\perp contains a totally isotropic sublattice of rank two). The Fourier expansion at a 1-dimensional cusp E is called Fourier–Jacobi expansion (see [4], [107], [40]). The value of a modular form G on the boundary component E is given by the Siegel operator $\Phi_E(G)$ (see [5]). This is the zeroth coefficient of the Fourier–Jacobi expansion which is a modular form with respect to a subgroup of $SL(2, \mathbb{Z})$.

The boundary of a 1-dimensional cusp E is a union of some 0-dimensional cusps. We consider the Fourier expansion of $\Phi_E(F|_{r^\perp})$ at a 0-dimensional cusp c associated to E as a part of the Fourier expansion of $(F|_{r^\perp})_{c,h}$ (see [74, Section 5.2] for the Fourier expansion of Fourier–Jacobi coefficients of modular forms). The indices of the Fourier coefficients of $\Phi_E(F|_{r^\perp})$ are of hyperbolic norm 0. Therefore $\Phi_E(F|_{r^\perp}) \equiv 0$ because, as shown above, all such coefficients in $(F|_{r^\perp})_{c,h}$ are equal to zero. Therefore the quasi pull-back $F|_{r^\perp}$ is a cusp form. \square

Using Theorem 8.18 we prove that the quasi pull-back defined in Theorem 8.3 is a cusp form.

Corollary 8.21. *Let $L \hookrightarrow \text{II}_{2,26}$ be a nondegenerate sublattice of signature $(2, n)$, $n \geq 1$. We assume that the set $R_{-2}(L^\perp)$ of (-2) -roots in L^\perp is non-empty. Then the quasi pull-back $\Phi|_L \in S_{12+N(L^\perp)}(\tilde{\mathcal{O}}(L), \det)$ of the Borcherds form Φ_{12} is a cusp form.*

Proof. To prove the corollary we divide the procedure of the quasi pull-back of Theorem 8.3 into finitely many steps.

First, we take a root $r_1 \in R_{-2}(L^\perp) \neq \emptyset$ and define $M_1 = (r_1)^\perp_{\text{II}_{2,26}}$. The form Φ_{12} has a zero of order 1 on $\mathcal{D}_{r_1}(\text{II}_{2,26})$. According to Theorem 8.18 we have

$$\Phi|_{M_1} = \frac{\Phi_{12}(Z)}{(r_1, Z)} \Big|_{M_1} \in S_{13}(\tilde{\mathcal{O}}^+(M_1), \det).$$

We note that the cusp form $\Phi|_{M_1}$ might have divisors different from the (-2) -divisors of Φ_{12} . If $s \in R_{-2}(L^\perp)$ such that $(r_1, s) \neq 0$ then $s_1 = \text{pr}_{M_1^\vee}(s)$ has negative norm $-2 < (s_1, s_1) < 0$ and $\Phi|_{M_1}$ vanishes along $\mathcal{D}_{s_1}(M_1)$ according to Lemma 8.2. Therefore the divisors of $\Phi|_{M_1}$ are rational quadratic divisors defined by some vectors in $v \in M_1^\vee$. But $\mathcal{D}_v(M_1) = \mathcal{D}_{\text{tv}}(M_1)$ and we can fix t in order to have a primitive vector $\text{tv} \in M_1$.

Second, we consider the lattice $L^\perp \cap M_1$. Suppose first that there is no vector $v \in L^\perp \cap M_1$ such that $\Phi|_{M_1}$ vanishes on $\mathcal{D}_v(M_1)$. In this case $\Phi|_L$ is equal to the pull-back of $\Phi|_{M_1}$ on \mathcal{D}_L . The pull-back of a cusp form is a cusp form and the proof is finished.

Otherwise, if $r_2 \in L^\perp \cap M_1$ is a vector such that $\Phi|_{M_1}$ vanishes on $\mathcal{D}_{r_2}(M_1)$, then we define $M_2 = (r_2)^\perp_{M_1} = (r_1, r_2)^\perp_{\text{II}_{2,26}}$. As in Theorem 8.18 we have

$$\Phi|_{M_2} = \frac{\Phi|_{M_1}}{(r_2, Z)^m} \Big|_{M_2} \in S_{13+m}(\tilde{\mathcal{O}}^+(M_2), \det)$$

where $m \geq 1$ is the degree of zero of $\Phi|_{M_1}$ on $\mathcal{D}_{r_2}(M_1)$.

The function $\Phi|_{M_2}$ is a modular form vanishing along some rational quadratic divisors. We can repeat the procedure described above for M_2 . After a finite number of steps we get the cusp form $\Phi|_L$. \square

Proposition 8.22. *Let L be one of the lattices L_{2d} , $L_{2,2d}$ or $L_{\mathbb{Q}(d)}$ defined in (7.10)–(7.14). We assume that there exists an embedding $L \hookrightarrow \text{II}_{2,26}$ such that the weight of the*

quasi pull-back $\Phi|_{\mathcal{L}}$ is smaller than the dimension of $\mathcal{D}_{\mathcal{L}}$. Then $\Phi|_{\mathcal{L}}$ vanishes along the ramification divisor of the modular projection

$$\pi: \mathcal{D}_{\mathcal{L}} \rightarrow \tilde{\mathcal{O}}^+(\mathcal{L}) \setminus \mathcal{D}_{\mathcal{L}}.$$

Proof. We give here a proof which works for all cases including the moduli spaces of polarised holomorphic symplectic O’Grady varieties (see [51, Corollary 4.6]).

The components of the branch divisor of π are $\mathcal{D}_r(\mathcal{L})$ where $r \in \mathcal{L}$ and σ_r or $-\sigma_r$ is in $\tilde{\mathcal{O}}^+(\mathcal{L})$ (see [46, Corollary 2.13] and Equation (6.12) above). If $\sigma_r \in \tilde{\mathcal{O}}^+(\mathcal{L})$, then $\Phi|_{\mathcal{L}}$ vanishes along $\mathcal{D}_r(\mathcal{L})$ because $\Phi|_{\mathcal{L}}$ is modular with character \det . We have to prove that $\Phi|_{\mathcal{L}}$ vanishes also on $\mathcal{D}_r(\mathcal{L})$ with $-\sigma_r \in \tilde{\mathcal{O}}^+(\mathcal{L})$. To prove this we use the transitivity of the quasi pull-back construction. Let $r \in \mathcal{L} \hookrightarrow \Pi_{2,26}$ and $L_r = r_{\mathcal{L}}^{\perp}$. Then

$$(\Phi|_{\mathcal{L}})|_{L_r} = \Phi|_{L_r}.$$

We have to consider three cases.

1) Let $r \in \mathcal{L} = L_{2d}$. Then $\text{rank } L_r = 18$, and, according to Corollary 7.19, $\det |(L_r)_{\Pi_{2,26}}^{\perp}|$ is either 1 or 4. In [23, Table I] one can find all classes of the indecomposable lattices of small rank and determinant. Analysing that table we find three classes of lattices of rank 8 of determinant 1, 2 or 4:

$$E_8 \ (|\mathcal{R}(E_8)| = 240), \quad E_7 \oplus A_1 \ (|\mathcal{R}(E_7 \oplus A_1)| = 128), \quad D_8 \ (|\mathcal{R}(D_8)| = 112).$$

Therefore $\Phi|_{12}$ has a zero of order at least 56 along $\mathcal{D}_r(\mathcal{L})$. The modular form $\Phi|_{\mathcal{L}}$ is of weight $k < 19$. Therefore $\mathcal{R}(L_{\Pi_{2,26}}^{\perp})$ has at most 12 roots. Therefore $\Phi|_{\mathcal{L}}$ vanishes on $\mathcal{D}_r(L_{2d})$ with order at least 50.

2) Let $r \in \mathcal{L} = L_{2,2d}$. Then $\text{rank } L_r = 19$. We described reflective vectors r with $-\sigma_r \in \tilde{\mathcal{O}}^+(\mathcal{L})$ in Proposition 7.18. There are three possible cases:

$$r^2 = 2d, \text{ div}(r) = d \text{ or } 2d \quad \text{and} \quad r^2 = d = \text{div}(r) \ (d \text{ is odd}).$$

According to Lemma 7.5, $\det L_r = 2, 4$ or 8 . Therefore the rank 7 lattice $(L_r)_{\Pi_{2,26}}^{\perp}$ has the same determinant. According to [23, Table I] there are six possible classes of such lattices:

$$E_7, \ D_7, \ D_6 \oplus A_1, \ A_7, \ [D_6 \oplus \langle 8 \rangle]_2, \ [E_6 \oplus \langle 24 \rangle]_3,$$

where $[M]_n$ denotes an overlattice of order n of M . Any of these lattices contains at least 60 roots ($|\mathcal{R}(D_6)| = 60$). The modular form $\Phi|_{\mathcal{L}}$ is of weight $k < 20$. Therefore $\mathcal{R}(L_{\Pi_{2,26}}^{\perp})$ has at most 14 roots and $\Phi|_{\mathcal{L}}$ vanishes on $\mathcal{D}_r(L_{2,2d})$ with order at least 23.

3) Let $r \in \mathcal{L} = L_{Q(d)}$. In this case $d \equiv 3 \pmod{4}$, and the discriminant group $D(L_{Q(d)})$ is cyclic of order d . Using Proposition 7.18 and Lemma 7.5 we obtain that only one class for $(L_r)_{\Pi_{2,26}}^{\perp}$ is possible. This is the lattice E_7 . We finish the proof as above. \square

9. Arithmetic of root systems

In this section we finish the proof of Theorem 6.1. To prove it we use the low weight cusp form trick (see the discussion in the proof of Theorem 6.13) by using the quasi pull-back of the Borcherds modular form Φ_{12} to construct cusp forms of small weight with large divisor.

According to Theorem 8.3, Theorem 8.18 and Proposition 8.22 the main point for us is the following. We want to know for which $2d > 0$ there exists a vector

$$(9.1) \quad l \in E_8, \quad l^2 = 2d, \quad l \text{ is orthogonal to at least 2 and at most 12 roots.}$$

To solve this problem we use the combinatorial geometry of the root system E_8 together with the theory of quadratic forms. First we give some properties of the lattice E_8 and we show how one can construct the first polarisations of general type in Theorem 6.1. After that we outline the answer to the question in (9.1).

9.1. Vectors in E_8 and E_7


By definition, the lattice D_8 is an even sublattice of the Euclidean lattice \mathbb{Z}^8

$$D_8 = \{l = (x_1, \dots, x_8) \in \mathbb{Z}^8 \mid x_1 + \dots + x_8 \in 2\mathbb{Z}\}.$$

The determinant of D_8 is equal to 4. We denote by e_1, \dots, e_8 the Euclidean basis of \mathbb{Z}^8 ($(e_i, e_j) = \delta_{ij}$). The lattice E_8 is the double extension of D_8 :

$$E_8 = D_8 \cup \left(\frac{e_1 + \dots + e_8}{2} + D_8 \right).$$

We consider the Coxeter basis of simple roots in E_8 (see [17])



where

$$\alpha_1 = \frac{1}{2}(e_1 + e_8) - \frac{1}{2}(e_2 + e_3 + e_4 + e_5 + e_6 + e_7),$$

$$\alpha_2 = e_1 + e_2, \quad \alpha_k = e_{k-1} - e_{k-2} \quad (3 \leq k \leq 8)$$

and $E_8 = \langle \alpha_1, \dots, \alpha_8 \rangle_{\mathbb{Z}}$. The lattice E_8 contains 240 roots. We recall that any root is a sum of simple roots with integral coefficients of the same sign. The fundamental weights ω_j of E_8 form the dual basis, so $(\alpha_i, \omega_j) = \delta_{ij}$. The formulae for the weights

are given in [17, Tabl. VII]. We shall use the Cartan matrix of the dual basis

$$(9.2) \quad ((\omega_i, \omega_j)) = \begin{pmatrix} 4 & 5 & 7 & 10 & 8 & 6 & 4 & 2 \\ 5 & 8 & 10 & 15 & 12 & 9 & 6 & 3 \\ 7 & 10 & 14 & 20 & 16 & 12 & 8 & 4 \\ 10 & 15 & 20 & 30 & 24 & 18 & 12 & 6 \\ 8 & 12 & 16 & 24 & 20 & 15 & 10 & 5 \\ 6 & 9 & 12 & 18 & 15 & 12 & 8 & 4 \\ 4 & 6 & 8 & 12 & 10 & 8 & 6 & 3 \\ 2 & 3 & 4 & 6 & 5 & 4 & 3 & 2 \end{pmatrix}.$$

Let us assume that $l_{2d} \in E_8$ and $(l_{2d})_{E_8}^\perp$ contains exactly 12 roots. We consider two cases

$$R((l_{2d})_{E_8}^\perp) = A_2 \oplus 3A_1 \quad \text{or} \quad R((l_{2d})_{E_8}^\perp) = A_2 \oplus A_2.$$

There are four possible choices of the subsystem $A_2 \oplus 3A_1$ inside the Dynkin diagram of E_8 . There are four choices for a copy of A_2 :

$$A_2 = \langle \alpha_1, \alpha_3 \rangle, \quad \langle \alpha_2, \alpha_4 \rangle, \quad \langle \alpha_5, \alpha_6 \rangle, \quad \langle \alpha_7, \alpha_8 \rangle.$$

Then the three pairwise orthogonal copies of A_1 are defined automatically. For example, if $A_2^{(1)} = \langle \alpha_5, \alpha_6 \rangle$ then $3A_1^{(1)} = \langle \alpha_2 \rangle \oplus \langle \alpha_3 \rangle \oplus \langle \alpha_8 \rangle$. The sum $A_2^{(1)} \oplus 3A_1^{(1)}$ is the root system of the orthogonal complement of the vector $l_{5,6} = \omega_1 + \omega_4 + \omega_7 \in E_8$. In fact, if $r = \sum_{i=1}^8 x_i \alpha_i$ is a positive root ($x_i \geq 0$) then $(r, l_{5,6}) = x_1 + x_4 + x_7 = 0$. Therefore $x_1 = x_4 = x_7 = 0$ and r belongs to $A_2^{(1)} \oplus 3A_1^{(1)}$ (see the Dynkin diagram of E_8 above). Using the matrix (9.2) we find $l_{5,6}^2 = 46$. Doing similar calculations with

$$l_{1,3} = \omega_4 + \omega_6 + \omega_8, \quad l_{2,4} = \omega_3 + \omega_5 + \omega_7, \quad l_{7,8} = \omega_1 + \omega_4 + \omega_6$$

we obtain polarisations for $d = 50, 54$ and 57

$$l_{1,3}^2 = 2 \cdot 50, \quad l_{2,4}^2 = 2 \cdot 54, \quad l_{7,8}^2 = 2 \cdot 57.$$

To get a good vector for $d = 52$ we consider the lattice $M = A_2 \oplus A_2 = \langle \alpha_3, \alpha_4 \rangle \oplus \langle \alpha_6, \alpha_7 \rangle$ in E_8 . Then M is the root system of the orthogonal complement of $l_{104} = \omega_1 + \omega_2 + \omega_5 + \omega_8$ and $l_{104}^2 = 2 \cdot 52$.

In this way we construct the first five polarisations of general type from the list of Theorem 6.1. These elementary arguments, similar to the arguments of Kondo in [76], do not give the whole list but only a few early cases. A sufficient condition for existence of vectors satisfying (9.1) is given in the theorem below.

Theorem 9.3. *A vector l satisfying (9.1) does exist if the inequality*

$$(9.4) \quad 4N_{E_7}(2d) > 28N_{E_6}(2d) + 63N_{D_6}(2d)$$

is valid, where $N_L(2d)$ denotes the number of representations of $2d$ by the lattice L .

Proof. We use bouquets of copies of A_2 in $E_8 \setminus E_7$. The root system $R(E_8)$ is a disjoint union of 126 roots of E_7 and the bouquet of 28 copies of A_2 centred in A_1 . This fact explains the coefficients 28 and 63 in the right hand side of (9.4). See [46, Section 7] for more details. \square

In [46] we found explicit formulae for the numbers of representations and we proved that

$$N_{E_7}(2m) > \frac{24\pi^3}{5\zeta(3)}, \quad 21m^2 > N_{D_6}(2d), \quad 103.69m^2 > N_{E_6}(2d).$$

These inequalities give a finite set of d for which (9.4) is not valid. Analysing the corresponding theta series we found the set of all such d , containing 131 numbers. The five tables in [46, Section 7] and the argument above for $d = 52$ give the list of polarisations of general type of Theorem 6.1.

The geometric genus of \mathcal{F}_{2d} is positive if there exists a cusp form of canonical weight 19. For each d not in the general type list but satisfying $d \geq 40$ and $d \neq 41, 44, 45$ or 47 , there exists a vector $h_{2d} \in E_8$ of length $2d$ orthogonal exactly to 14 roots. The corresponding quasi pull-back is a cusp forms of canonical weight. For $d = 42, 43, 51, 53$ and 55 such cusp forms were constructed by Kondo in [76]. For other d see [46].

A similar arithmetic method applied to E_7 (see [50, Section 4]) gives the proof of Theorem 6.2. There is a significant technical difficulty in the case of E_7 . The proof involves estimating the number of ways of representing certain integers by various root lattices of odd rank. In [50, Section 5]) we gave a new, clear, explicit version of Siegel's formula for this number in the odd rank case. It may be expressed either in terms of Zagier L-functions or in terms of the H. Cohen numbers. For example we obtained a new, very short formula for the number of representations by 5 squares (see [50, Section 4] for the details).

9.2. Binary quadratic forms in E_8

We saw in Example 4.5 that the results of Debarre and Voisin [25] imply that the moduli space $\mathcal{M}_{2,11}^{[2],\text{non-split}}$ of Beauville degree $d = 11$ is unirational. We note that $\mathcal{M}_{2,11}^{[2],\text{split}}$, which is a finite covering of $\mathcal{M}_{2,11}^{[2],\text{non-split}}$ by (7.16), has non-negative Kodaira dimension. Theorem 6.2 shows that there can be at most 11 exceptional split polarisations of non general type. Proposition 9.5 hints that one can expect a theorem for the non-split case, which we hope to prove in the future, similar to Theorem 6.2.

We recall that the Beauville degree $d \equiv -1 \pmod{4}$ if the polarisation h_{2d} is of non-split type. For small d we can calculate the class of the orthogonal complement of $Q(d)$. According to [23, Table 0] the rank 6 lattice $Q(d)_{E_8}^\perp$ of determinant d contains at least 24 roots for $d = 3, 7, 11$ and 15 . One can continue this analysis but

we propose below a simple algorithm which gives us the first “good” embeddings of $Q(d)$ in E_8 .

Proposition 9.5. *The moduli spaces $\mathcal{M}_{2,39}^{[2],non-split}$ and $\mathcal{M}_{2,47}^{[2],non-split}$ are of general type.*

Proof. Let $Q(d)(-1) = \begin{pmatrix} 2 & -1 \\ -1 & 2c \end{pmatrix}$, where $c = (d + 1)/4$, be the binary quadratic form associated to a non-split polarisation of degree $2d$ (see (7.14)). To make the notation simpler we denote this binary form by $Q(d)$. We have to embed $Q(d)$ in E_8 so as to satisfy

$$2 \leq |R(Q(d) \frac{1}{E_8})| \leq 14.$$

We describe below a method (we call it $\frac{1}{2}(++)$ -algorithm) which gives many such embeddings. We take the following realisation of the binary quadratic form of determinant d in E_8 :

$$Q(d) = \langle e_2 - e_1, v_{2c} \rangle = \langle e_2 - e_1, \frac{1}{2}(e_1 - e_2 + x_3 e_3 + \dots + x_8 e_8) \rangle,$$

where $x_i \geq 0$ and $2c = (2 + x_3^2 + \dots + x_8^2)/4$. The second generator v_{2c} of the binary quadratic form is a half-integral element of E_8 . According to the definition of E_8 given above

$$v_{2c} + \frac{1}{2}(e_1 + \dots + e_8) = e_1 + \frac{x_3 + 1}{2} e_3 + \dots + \frac{x_8 + 1}{2} e_8 \in D_8.$$

Therefore $x_3 \equiv \dots \equiv x_8 \pmod{2}$ and $x_3 + \dots + x_8 \equiv 0 \pmod{4}$. Now we can find all roots orthogonal to $Q(d)$.

First, we consider the integral roots of E_8 . The roots $\pm(e_1 + e_2)$ are orthogonal to $Q(d)$ therefore $R(Q(d) \frac{1}{E_8})$ is not empty. Then the integral roots $\pm(e_i \pm e_j)$ ($i, j > 2$) are orthogonal to $Q(d)$ if and only if $x_i = \mp x_j$.

Second, a half-integral root orthogonal to $Q(d)$ has the form

$$\pm \frac{1}{2}(e_1 + e_2 + \sum_{i=3}^8 (-1)^{\epsilon_i} e_i)$$

where the number of $-$ signs in the sum is even. Therefore v_{2c} is orthogonal to a half-integral root if and only if we can divide the vector of coefficients (x_3, \dots, x_8) in two parts containing even number of terms and with the same sum. If this is possible there are two pairs of roots orthogonal to v_{2c} . For example, if there is a root of type $\pm(++; ++++--)$ then we also have $\pm(++; ----++)$. The rules described above give the number (it is always positive) of roots orthogonal to $Q(d)$.

To finish the proof of Proposition 9.5 we consider two vectors

$$v_{20} = \frac{1}{2}(1, 1; 5, 5, 3, 3, 3, 1) \quad \text{and} \quad v_{24} = \frac{1}{2}(1, 1; 7, 5, 3, 3, 1, 1).$$

The quadratic forms $Q(39)$ and $Q(47)$ are orthogonal to exactly 14 roots in E_8 . As above, using Theorem 8.3, Theorem 8.18 and Proposition 8.22 we see that the quasi pull-back of Φ_{12} on the modular variety defined by the lattice $L_{Q(d)}$ is a cusp

form of weight 19 vanishing on the ramification divisor. We finish the proof using Theorem 6.13. \square

References

- [1] A. Apostolov. *Moduli spaces of polarized irreducible symplectic manifolds are not necessarily connected*. arXiv:1109.0175, 17 pp. \leftarrow [480](#)
- [2] M. Artin. Supersingular K3 surfaces. *Ann. Scient. Éc. Norm. Sup. (4)*, 7, 543–568, 1974. \leftarrow [462](#)
- [3] A. Ash, D. Mumford, M. Rapoport, and Y. Tai. *Smooth compactification of locally symmetric varieties*. Second edition: Cambridge University Press, Cambridge, 2010. \leftarrow [487](#), [488](#), [489](#), [494](#), [498](#)
- [4] W. L. Baily. Fourier-Jacobi series, in: Algebraic Groups and Discontinuous Subgroups, 296–300. *Proc. Sympos. Pure Math.*, 9, Amer. Math. Soc., Providence, 1966. \leftarrow [496](#), [511](#)
- [5] W. L. Baily Jr. and A. Borel. Compactification of arithmetic quotients of bounded symmetric domains. *Ann. of Math. (2)*, 84, 442–528, 1966. \leftarrow [469](#), [483](#), [485](#), [486](#), [511](#)
- [6] W. Barth, K. Hulek, C. Peters, and A. Van de Ven. Compact complex surfaces. Second Edition. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, 4, Springer, Berlin, 2004. \leftarrow [461](#), [463](#), [464](#), [465](#), [466](#), [468](#)
- [7] J. Bauermann. Lokale Tensorfelder auf arithmetischen Quotienten symmetrischer beschränkter Gebiete. *J. Reine Angew. Math.*, 428, 111–138, 1992. \leftarrow [496](#)
- [8] A. Beauville. Variétés Kähleriennes dont la première classe de Chern est nulle. *J. Differential Geom.*, 18, 755–782, 1983. \leftarrow [462](#), [473](#), [474](#), [475](#)
- [9] F. Bogomolov. The decomposition of Kähler manifolds with a trivial canonical class. *Mat. Sb. (N.S.)*, 93(135), 573–575, 630, 1974. \leftarrow [462](#), [473](#), [475](#)
- [10] F. Bogomolov. Hamiltonian Kählerian manifolds. *Dokl. Akad. Nauk SSSR*, 243, 1101–1104, 1978. \leftarrow [473](#)
- [11] R. E. Borcherds. Automorphic forms on $O_{s+2,2}(\mathbb{R})$ and infinite products. *Invent. Math.*, 120, 161–213, 1995. \leftarrow [495](#), [499](#), [500](#), [505](#), [506](#)
- [12] R. E. Borcherds. The moduli space of Enriques surfaces and the fake monster Lie superalgebra. *Topology*, 35, 699–710, 1996. \leftarrow [508](#)
- [13] R. E. Borcherds. Automorphic forms with singularities on Grassmannians. *Invent. Math.*, 132, 491–562, 1998. \leftarrow [497](#)
- [14] R. E. Borcherds, L. Katzarkov, T. Pantev, and N. I. Shepherd-Barron. Families of K3 surfaces. *J. Algebraic Geom.*, 7, 183–193, 1998. \leftarrow [505](#), [506](#)
- [15] A. Borel. Some metric properties of arithmetic quotients of symmetric spaces and an extension theorem. *J. Differential Geom.*, 6, 543–560, 1972. \leftarrow [472](#)
- [16] A. Borel and L. Ji. Compactifications of symmetric and locally symmetric spaces. *Mathematics: Theory & Applications*, Birkhäuser, Boston, 2006. \leftarrow [484](#), [485](#), [486](#)

- [17] N. Bourbaki. *Groupes et Algèbres de Lie, Chapitres 4, 5 et 6*. Hermann, Paris, 1968. ← [514](#), [515](#)
- [18] R. Brussee. The canonical class and the C^∞ properties of Kähler surfaces. *New York J. Math.*, **2**, 103–146, 1996. ← [461](#)
- [19] N. Buchdahl. On compact Kähler surfaces. *Ann. Inst. Fourier (Grenoble)*, **49**, 287–302, 1999. ← [461](#)
- [20] D. Burns and M. Rapoport. On the Torelli problem for Kählerian K3 surfaces. *Ann. Sc. ENS.*, **8**, 235–274, 1975. ← [461](#), [467](#)
- [21] H. Cartan. Quotient d’un espace analytique par un groupe d’automorphismes, in: Algebraic geometry and topology, a symposium in honor of S. Lefschetz, 90–102. Princeton University Press, Princeton, 1957. ← [490](#)
- [22] C. Chevalley. Invariants of finite groups generated by reflections. *Amer. J. Math.*, **77**, 778–782, 1955. ← [483](#)
- [23] J. H. Conway and N. J. A. Sloane. Low-dimensional lattices I: quadratic forms of small determinant. *Proc. Roy. Soc. London Ser. A*, **418**(1854), 17–41, 1988. ← [513](#), [516](#)
- [24] O. Debarre. Un contre-exemple au théorème de Torelli pour les variétés symplectiques irréductibles. *C. R. Acad. Sci. Paris Sér. I Math.*, **299**, 681–684, 1984. ← [462](#), [475](#)
- [25] O. Debarre and C. Voisin. Hyper-Kähler fourfolds and Grassmann geometry. *J. Reine Angew. Math.*, **649**, 63–87, 2010. ← [482](#), [516](#)
- [26] I. Dolgachev. Mirror symmetry for lattice polarized K3 surfaces. *J. Math. Sci.*, **81**, 2599–2630, 1996. ← [471](#)
- [27] I. Dolgachev, B. van Geemen, and S. Kondō. A complex ball uniformization of the moduli space of cubic surfaces via periods of K3 surfaces. *J. Reine Angew. Math.*, **588**, 99–148, 2005. ← [481](#)
- [28] I. Dolgachev and S. Kondō. Moduli of K3 surfaces and complex ball quotients, in: Arithmetic and geometry around hypergeometric functions, 43–100. *Progr. Math.*, **260**, Birkhäuser, Basel, 2007. ← [481](#)
- [29] M. Dürr. *Seiberg-Witten theory and the C^∞ -classification of complex surfaces*. Ph.D. thesis, Zürich, 2002. ← [461](#)
- [30] M. Eichler. Quadratische Formen und orthogonale Gruppen. *Grundlehren der mathematischen Wissenschaften*, **63**, Springer, Berlin–New York, 1974. ← [503](#), [509](#)
- [31] G. Faltings and C.-L. Chai. Degeneration of abelian varieties. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, **22**. Springer, Berlin, 1990. ← [490](#), [494](#)
- [32] E. Freitag. Siegelische Modulfunktionen. *Grundlehren der mathematischen Wissenschaften*, **254**. Springer, Berlin–Göttingen–Heidelberg, 1983. ← [496](#), [498](#)
- [33] E. Freitag. Some modular forms related to cubic surfaces. *Kyungpook Math. J.*, **43**, 433–462, 2003. ← [508](#)

- [34] R. Friedman. A new proof of the global Torelli theorem for K3 surfaces. *Ann. of Math. (2)*, **120**, 237–269, 1984. ← [461](#), [482](#)
- [35] R. Friedman and J. W. Morgan. Algebraic surfaces and Seiberg-Witten invariants. *J. Algebraic Geom.*, **6**, 445–479, 1997. ← [461](#)
- [36] A. Fujiki. On the de Rham cohomology group of a compact Kähler symplectic manifold, in: Algebraic geometry (Sendai 1985), 105–165. *Adv. Stud. Pure Math.*, **10**, North-Holland, Amsterdam, 1987. ← [474](#)
- [37] A. Garbagnati and A. Sarti. Elliptic fibrations and symplectic automorphisms on K3 surfaces. *Comm. Algebra*, **37**, 3601–3631, 2009. ← [481](#)
- [38] V. Gritsenko. Jacobi functions of n -variables. *Zap. Nauk. Sem. LOMI*, **168**, 32–45, 1988. English translation in *J. Soviet Math.*, **53**, 243–252, 1991. ← [496](#)
- [39] V. Gritsenko. Irrationality of the moduli spaces of polarized Abelian surfaces. *Internat. Math. Res. Notices*, **6**, 235–243, 1994. ← [496](#)
- [40] V. Gritsenko. Modular forms and moduli spaces of Abelian and K3 surfaces. *Algebra i Analiz*, **6**, 65–102, 1994; English translation in: *St. Petersburg Math. J.*, **6**, 1179–1208, 1995. ← [495](#), [496](#), [498](#), [503](#), [511](#)
- [41] V. Gritsenko. Reflective modular forms in algebraic geometry, 28 pp. [arXiv:1005.3753](#) ← [500](#), [501](#)
- [42] V. Gritsenko and K. Hulek. Appendix to the paper “Irrationality of the moduli spaces of polarized abelian surfaces”. Abelian varieties, Egloffstein 1993, 83–84. Walter de Gruyter, Berlin, 1995. ← [498](#)
- [43] V. Gritsenko and K. Hulek. Minimal Siegel modular threefolds. *Math. Proc. Cambridge Philos. Soc.*, **123**, 461–485, 1998. ← [476](#), [496](#)
- [44] V. Gritsenko and K. Hulek. Commutator coverings of Siegel threefolds. *Duke Math. J.*, **94**, 509–542, 1998. ← [496](#)
- [45] V. Gritsenko and K. Hulek. Uniruledness of orthogonal modular varieties. 14 pp., [arXiv:1202.3361](#). ← [501](#)
- [46] V. Gritsenko, K. Hulek, and G. K. Sankaran. The Kodaira dimension of the moduli of K3 surfaces. *Invent. Math.*, **169**, 519–567, 2007. ← [461](#), [490](#), [493](#), [494](#), [495](#), [497](#), [504](#), [505](#), [507](#), [513](#), [516](#)
- [47] V. Gritsenko, K. Hulek, and G. K. Sankaran. The Hirzebruch-Mumford volume for the orthogonal group and applications. *Documenta Math.*, **12**, 215–241, 2007. ← [504](#)
- [48] V. Gritsenko, K. Hulek, and G. K. Sankaran. Hirzebruch-Mumford proportionality and locally symmetric varieties of orthogonal type. *Documenta Math.*, **13**, 1–19, 2008. ← [470](#), [499](#)
- [49] V. Gritsenko, K. Hulek, and G. K. Sankaran. Abelianisation of orthogonal groups and the fundamental group of modular varieties. *J. of Algebra*, **322**, 463–478, 2009. ← [469](#), [496](#), [500](#), [503](#), [509](#)
- [50] V. Gritsenko, K. Hulek, and G. K. Sankaran. Moduli spaces of irreducible symplectic manifolds. *Compositio Math.*, **146**, 404–434, 2010. ← [462](#), [472](#),

[477](#), [478](#), [479](#), [480](#), [495](#), [502](#), [504](#), [505](#), [516](#)

- [51] V. Gritsenko, K. Hulek, and G. K. Sankaran. Moduli spaces of polarised symplectic O'Grady varieties and Borchers products. *J. Differential Geom.*, **88**, 61–85, 2011. ← [462](#), [478](#), [495](#), [504](#), [505](#), [507](#), [513](#)
- [52] V. Gritsenko and V. Nikulin. K3 surfaces, Lorentzian Kac–Moody algebras and mirror symmetry. *Math. Res. Lett.*, **3**, 211–229, 1996. ← [471](#)
- [53] V. Gritsenko and V. Nikulin. Siegel automorphic form correction of some Lorentzian Kac–Moody Lie algebras. *Amer. J. Math.*, **119**, 181–224, 1997. ← [496](#)
- [54] V. Gritsenko and V. Nikulin. Automorphic forms and Lorentzian Kac–Moody algebras. I. *International J. Math.*, **9**(2), 153–200, 1998. ← [499](#)
- [55] V. Gritsenko and V. Nikulin. Automorphic forms and Lorentzian Kac–Moody algebras. II. *International J. Math.*, **9** (2), 201–275, 1998. ← [496](#), [497](#), [507](#), [508](#), [509](#)
- [56] V. Gritsenko and V. Nikulin. The arithmetic mirror symmetry and Calabi–Yau manifolds. *Comm. Math. Phys.*, **200**, 1–11, 2000. ← [471](#)
- [57] V. Gritsenko and G. K. Sankaran. Moduli of abelian surfaces with a $(1, p^2)$ polarisation. *Izv. Ross. Akad. Nauk Ser. Mat.*, **60**, 19–26, 1996 ; reprinted in: *Izv. Math.*, **60**, 893–900, 1996. ← [498](#)
- [58] B. Hassett and Y. Tschinkel. Moving and ample cones of holomorphic symplectic fourfolds. *Geometric and Functional Analysis*, **19**, 1065–1080, 2009. ← [478](#)
- [59] B. Hassett and Y. Tschinkel. Flops on holomorphic symplectic fourfolds and determinantal cubic hypersurfaces. *J. Inst. Math. Jussieu*, **9**, 125–153, 2010. ← [478](#)
- [60] S. Helgason. Differential geometry, Lie groups, and symmetric spaces. *Graduate Studies in Mathematics*, **34**, Amer. Math. Soc., Providence, 2001. ← [485](#)
- [61] E. Horikawa. Surjectivity of the period map of K3 surfaces of degree 2. *Math. Ann.*, **228**, 113–146, 1977. ← [461](#)
- [62] E. Horikawa. On the periods of Enriques surfaces I. *Proc. Japan Acad.*, **53**, 124–127, 1977. ← [481](#)
- [63] D. Huybrechts. Compact hyper-Kähler manifolds: basic results. *Invent. Math.*, **135**, 63–113, 1999. ← [462](#), [475](#)
- [64] D. Huybrechts. A global Torelli theorem for hyperkähler manifolds [after Verbitsky]. Séminaire BOURBAKI 63ème année, n 1040, 2010–2011. ← [462](#), [476](#)
- [65] A. Iliev and K. Ranestad. K3 surfaces of genus 8 and varieties of sums of powers of cubic fourfolds. *Trans. Amer. Math. Soc.*, **353**, 1455–1468, 2001. ← [482](#)
- [66] A. Iliev and K. Ranestad. Addendum to K3 surfaces of genus 8 and varieties of sums of powers of cubic fourfolds. *C. R. Acad. Bulgare Sci.*, **60**, 1265–1270, 2007. ← [482](#)
- [67] D. Kaledin, M. Lehn, and C. Sorger. Singular symplectic moduli spaces. *Invent. Math.*, **164**, 591–614, 2006. ← [473](#)

- [68] Y. Kawamata. Unobstructed deformations. A remark on a paper of Z. Ran. *J. Algebraic Geom.*, **1**, 183–190, 1992. ← [474](#)
- [69] D. Kazhdan. On the connection of the dual space of a group with the structure of its closed subgroups. *Funkcional. Anal. i Pril.*, **1**, 71–74, 1967. ← [496](#)
- [70] M. Kneser. Erzeugung ganzzahliger orthogonaler Gruppen durch Spiegelungen. *Math. Ann.*, **255**, 453–462, 1981. ← [480](#)
- [71] K. Kodaira. On the structure of compact complex analytic surfaces. I. *Amer. J. Math.*, **86**, 751–789, 1964. ← [461](#)
- [72] K. Kodaira and D. C. Spencer. On deformations of complex analytic structures. III. Stability theorems for complex structures. *Ann. of Math. (2)*, **71**, 43–76, 1960. ← [467](#)
- [73] J. Kollár and T. Matsusaka. Riemann-Roch type inequalities. *Amer. J. Math.*, **105**, 229–252, 1983. ← [478](#)
- [74] S. Kondo. On the Kodaira dimension of the moduli space of K3 surfaces. *Compositio Math.*, **89**, 251–299, 1993. ← [461](#), [494](#), [512](#)
- [75] S. Kondo. The rationality of the moduli space of Enriques surfaces. *Compositio Math.*, **91**, 159–173, 1994. ← [481](#)
- [76] S. Kondo. On the Kodaira dimension of the moduli space of K3 surfaces. II. *Compositio Math.*, **116**, 111–117, 1999. ← [505](#), [515](#), [516](#)
- [77] V. Kulikov. Degenerations of K3 surfaces and Enriques surfaces. *Izv. Akad. Nauk SSSR Ser. Mat.*, **41**, 1008–1042, 1199, 1977. ← [461](#)
- [78] A. Lamari. Courants kählériens et surfaces compactes. *Ann. Inst. Fourier (Grenoble)*, **49**, 263–285, 1999. ← [461](#)
- [79] D. Lieberman and D. Mumford. Matsusaka’s big theorem, in: Algebraic Geometry, Arcata 1974, 513–530. *Proc. Sympos. Pure Math.*, **29**, Amer. Math. Soc., Providence, 1975. ← [472](#)
- [80] E. Looijenga. A Torelli theorem for Kähler-Einstein K3 surfaces. *Lecture Notes in Math.*, **894**, 107–112. Springer, Berlin–New York, 1981. ← [461](#)
- [81] E. Looijenga and C. Peters. Torelli theorems for Kähler K3 surfaces. *Compositio Math.*, **42**, 145–186, 1981. ← [461](#)
- [82] S. Ma. The unirationality of the moduli spaces of 2-elementary K3 surfaces. With an appendix by K. Yoshikawa. *Proc. London Math. Soc.*, doi: 10.1112/plms/pds008, 2012. ← [481](#)
- [83] E. Markman. On the monodromy of moduli spaces of sheaves on K3 surfaces II, 76 pp. [math.AG/0305043](#) ← [476](#), [479](#)
- [84] E. Markman. On the monodromy of moduli spaces of sheaves on K3 surfaces. *J. Algebraic Geom.*, **17**, 29–99, 2008. ← [476](#)
- [85] E. Markman. Integral constraints on the monodromy group of the hyperkahler resolution of a symmetric product of a K3 surface. *Internat. J. Math.*, **21**, 169–223, 2010. ← [476](#)

- [86] E. Markman. A survey of Torelli and monodromy results for holomorphic-symplectic varieties, in: *Complex and Differential Geometry, Hannover 2009*, 257–322. *Springer Proceedings in Mathematics*, 8, Springer, Berlin, 2011. ← [462](#), [476](#), [477](#), [478](#), [479](#)
- [87] T. Matsusaka. Polarized varieties with a given Hilbert polynomial. *Amer. J. Math.*, 94, 1027–1077, 1972. ← [472](#), [478](#)
- [88] D. Morrison. Some remarks on the moduli of K3 surfaces, in: *Classification of algebraic and analytic manifolds*, 303–332. *Progr. Math.*, 39, Birkhäuser, Boston, 1983. ← [470](#)
- [89] S. Mukai. Curves, K3 surfaces and Fano 3-folds of genus ≤ 10 , in: *Algebraic geometry and commutative algebra*, Vol. I, 357–377. Kinokuniya, Tokyo, 1988. ← [461](#), [481](#)
- [90] S. Mukai. Polarized K3 surfaces of genus 18 and 20, in: *Complex projective geometry, Trieste 1989/Bergen 1989*, 264–276. *London Math. Soc. Lecture Note Ser.*, 179, Cambridge Univ. Press, Cambridge, 1992. ← [461](#), [481](#)
- [91] S. Mukai. Curves and K3 surfaces of genus eleven, in: *Moduli of vector bundles, Sanda 1994/Kyoto 1994*, 189–197. *Lecture Notes in Pure and Appl. Math.*, 179, Dekker, New York, 1996. ← [461](#), [481](#)
- [92] S. Mukai. Polarized K3 surfaces of genus thirteen, in: *Moduli spaces and arithmetic geometry, Kyoto 2004*, 315–326, *Adv. Stud. Pure Math.*, 45, Math. Soc. Japan, Tokyo, 2006. ← [461](#), [481](#)
- [93] S. Mukai. Polarized K3 surfaces of genus sixteen. Oberwolfach Report 02/2010. DOI: 10.4171/OWR/2010/02 ← [461](#), [481](#)
- [94] D. Mumford. Hirzebruch’s proportionality theorem in the noncompact case. *Invent. Math.*, 42, 239–272, 1977. ← [499](#)
- [95] Y. Namikawa. Counter-example to global Torelli problem for irreducible symplectic manifolds. *Math. Ann.*, 324, 841–845, 2002. ← [462](#), [476](#)
- [96] V. V. Nikulin. Finite automorphism groups of Kähler K3 surfaces. *Trudy Moskov. Mat. Obshch.*, 38, 75–137, 1979; English translation in *Trans. Mosc. Math. Soc.*, 2, 71–135, 1980. ← [471](#)
- [97] V. V. Nikulin. Integral symmetric bilinear forms and some of their applications. *Izv. Akad. Nauk SSSR Ser. Mat.*, 43, 111–177, 1979; English translation in *Math. USSR, Izvestiia*, 14, 103–167, 1980. ← [469](#), [502](#), [506](#)
- [98] N. O. Nygaard. The Torelli theorem for ordinary K3 surfaces over finite fields, in: *Arithmetic and geometry, Volume I. Progr. Math.*, 36, 267–276, Birkhäuser, Boston 1983. ← [462](#)
- [99] K. O’Grady. Desingularized moduli spaces of sheaves on a K3. *J. Reine Angew. Math.*, 512, 49–117, 1999. ← [462](#), [473](#), [495](#)
- [100] K. O’Grady. A new six-dimensional irreducible symplectic variety. *J. Algebraic Geom.*, 12, 435–505, 2003. ← [462](#), [473](#)

- [101] K. O’Grady. Irreducible symplectic 4-folds and Eisenbud-Popescu-Walter sextics. *Duke Math. J.*, **134**, 99–137, 2006. ← [482](#)
- [102] A. Ogus. A crystalline Torelli theorem for supersingular K3 surfaces, in: Arithmetic and geometry, Volume II, 361–394. *Progr. Math.*, **36**, Birkhäuser, Boston 1983. ← [462](#)
- [103] C. Okonek and A. Van de Ven. Stable bundles, instantons and C^∞ -structures on algebraic surfaces, in: Several complex variables VI, 197–249. *Encyclopaedia Math. Sci.*, **69**, Springer, Berlin, 1990. ← [461](#)
- [104] M. Olsson. Semistable degenerations and period spaces for polarized K3 surfaces. *Duke Math. J.*, **125**, 121–203, 2004. ← [482](#)
- [105] U. Persson and H. Pinkham. Degeneration of surfaces with trivial canonical bundle. *Ann. of Math. (2)*, **113**, 45–66, 1981. ← [461](#)
- [106] A. Peterson and G. K. Sankaran. On some lattice computations related to moduli problems. With an appendix by V. Gritsenko. *Rend. Semin. Mat. Univ. Politec. Torino*, **68**, 289–304, 2010. ← [462](#)
- [107] I. Piatetskii-Shapiro. *Automorphic functions and the geometry of classical domains*. Gordon and Breach, New York, 1969. ← [496](#), [511](#)
- [108] I. Piatetskii-Shapiro and I. Shafarevich. A Torelli theorem for algebraic surfaces of type K3. *Izv. Akad. Nauk SSSR, Ser. Mat.*, **35**, 530–572, 1971; English translation in: *Math. USSR, Izv.*, **5** (1971), 547–588, 1972. ← [461](#)
- [109] Z. Ran. Deformations of manifolds with torsion or negative canonical bundle. *J. Algebraic Geom.*, **1**, , 279–291, 1992. ← [474](#)
- [110] A. Rapagnetta. Topological invariants of O’Grady’s six dimensional irreducible symplectic variety. *Math. Z.*, **256**, 1–34, 2007. ← [474](#)
- [111] A. Rapagnetta. On the Beauville form of the known irreducible symplectic varieties. *Math. Ann.*, **340**, 77–95, 2008. ← [474](#)
- [112] M. Reid. Young person’s guide to canonical singularities, in: Algebraic geometry, Bowdoin 1985, 345–414. *Proc. Sympos. Pure Math.*, **46**, Part 1, Amer. Math. Soc., Providence, 1987. ← [490](#)
- [113] B. Saint-Donat. Projective models of K3 surfaces. *Amer. J. Math.*, **96**, 602–639, 1974. ← [468](#), [472](#)
- [114] I. Satake. Algebraic structures of symmetric domains. *Kano Memorial Lectures*, **4**, Iwanami Shoten, Tokyo; Princeton University Press, Princeton, 1980. ← [469](#), [486](#)
- [115] F. Scattone. On the compactification of moduli spaces of algebraic K3 surfaces. *Mem. Amer. Math. Soc.*, **70**(374), 1987. ← [482](#), [494](#)
- [116] M. Schütt. Arithmetic of K3 surfaces. *Jahresber. Deutsch. Math.-Verein.*, **111**, 23–41, 2009. ← [462](#)
- [117] J. Shah. Surjectivity of the period map in the case of quartic surfaces and sextic double planes. *Bull. Amer. Math. Soc.*, **82**, 716–718, 1976. ← [461](#)

- [118] J. Shah. A complete moduli space for K3 surfaces of degree 2. *Ann. of Math.* (2), **112**, 485–510, 1980. ← [461](#), [482](#)
- [119] J. Shah. Degenerations of K3 surfaces of degree 4. *Trans. Amer. Math. Soc.*, **263**, 271–308, 1981. ← [461](#), [482](#)
- [120] N. I. Shepherd-Barron. The rationality of some moduli spaces of plane curves. *Compositio Math.*, **67**, 51–88, 1988. ← [481](#)
- [121] N. I. Shepherd-Barron. Apolarity and its applications. *Invent. Math.*, **97**, 433–444, 1989. ← [481](#)
- [122] Y. T. Siu. Every K3 surface is Kähler. *Invent. Math.*, **73**, 139–150, 1983. ← [461](#), [463](#)
- [123] H. Sterk. Lattices and K3 surfaces of degree 6. *Linear Algebra Appl.*, **226/228**, 297–309, 1995. ← [482](#)
- [124] Y. Tai. On the Kodaira dimension of the moduli space of abelian varieties. *Invent. Math.*, **68**, 425–439, 1982. ← [491](#)
- [125] G. Tian. Smoothness of the universal deformation space of compact Calabi-Yau manifolds and its Petersson-Weil metric, in: Mathematical aspects of string theory, 629–646. *Advanced Series in Mathematical Physics*, **1**, World Scientific, Singapore, 1987. ← [474](#)
- [126] A. N. Todorov. The Weil-Petersson geometry of the moduli space of $SU(n \geq 3)$ (Calabi-Yau) manifolds I. *Comm. Math. Phys.*, **126**, 325–346, 1989. ← [461](#), [468](#), [474](#)
- [127] M. Verbitsky. A global Torelli theorem for hyperkahler manifolds , 46 pp. arXiv:0908.4121v7 ← [462](#), [476](#), [477](#), [478](#)
- [128] E. Viehweg. Quasi-projective moduli for polarized manifolds. *Ergebnisse der Mathematik und ihrer Grenzgebiete* (3), **30**, Springer, Berlin, 1995. ← [471](#)
- [129] C. Voisin. Théorème de Torelli pour les cubiques de \mathbb{P}^5 . *Invent. Math.*, **86**, 577–601, 1986. ← [462](#), [482](#)
- [130] C. Voisin. *Géométrie des espaces de modules de courbes et de surfaces K3 [d'après Gritsenko-Hulek-Sankaran, Farkas-Popa, Mukai, Verra...]*. Séminaire BOURBAKI 59ème année, 2006–2007, n 981. ← [481](#)
- [131] A. Weil. *Oeuvres scientifiques – Collected papers*, Volumes I, II, III, Springer, New York–Heidelberg, 1979. ← [460](#)
- [132] M. Zowislok. On moduli spaces of sheaves on K3 or abelian surfaces, 24 pp. arXiv:1104.5629, ← [473](#)

Université Lille 1, Laboratoire Paul Painlevé, F-59655 Villeneuve d'Ascq, Cedex, France
 E-mail address: Valery.Gritsenko@agat.univ-lille1.fr

Institut für Algebraische Geometrie, Leibniz Universität Hannover, D-30060 Hannover, Germany
 E-mail address: hulek@math.uni-hannover.de

Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, England
 E-mail address: G.K.Sankaran@bath.ac.uk

